

---

# I Wish Someone Had Told Me This About Self Harm

---

## I Wish Someone Had Told Me This About

*Self-Harm Understanding what can sit underneath the part other people see. Self-Harm is best understood in context. In this guide we are looking at a behaviour that can have different functions and meanings and deserves understanding without punishment. That matters because short descriptions can make people compare themselves with a stereotype and conclude that their experience does not count. Real life is usually messier: symptoms overlap, people mask, responsibilities continue, and somebody can function in one setting while paying heavily for it later. Common patterns can include overwhelming emotion, numbness, dissociation, self-directed anger, communication of distress or attempts to regain control. None of these behaviours should be read in isolation. The same outward behaviour can have very different meanings for different people. The useful question is not "Why are you doing this?" in an accusing sense, but "What is this response trying to achieve, prevent, escape or make more manageable?" The body and mind are not separate systems. Sleep, pain, sensory load, hormones, physical illness, medication, grief, trauma, neurodivergence, financial pressure, caring responsibilities and relationships can all change how distress is experienced. A good assessment therefore looks beyond a checklist and asks what has changed, what the person is carrying and what makes the difficulty better or worse. Many people become skilled at hiding distress. They may smile, work, care for children, attend appointments and answer messages while privately struggling. Functioning should not be confused with being unaffected. Sometimes the cost appears afterwards as shutdown, tears, irritability, exhaustion, withdrawal or a need for long recovery time. Understanding self-harm also means noticing the cost of coping. People may be using most of their available energy to appear fine, complete essential tasks or prevent others from worrying. When that effort is invisible, support can arrive too late because everyone assumes that continued functioning means continued capacity. A useful starting point is to replace judgement with observation: what has changed, what is becoming harder, what is being avoided, what requires more recovery, and what does the person say they are frightened of or trying to manage? Those questions create a fuller picture than asking whether they meet somebody else's idea of what this condition should look like.*

---

## What people often misunderstand

One misunderstanding is that self-harm should always look obvious. In reality, people adapt. They may build routines, avoidance strategies, scripts, checking habits or ways of appearing composed. Those adaptations can keep life moving, but they can also hide how restricted or exhausted the person has become. Another misunderstanding is that insight should automatically switch a symptom off. Knowing intellectually that a response is unhelpful is not the same as being able to stop a nervous-system pattern, compulsion, trauma response or entrenched coping strategy on command. Change usually requires safety, repetition, support and a realistic pace. Language matters. "Attention seeking", "dramatic", "lazy", "controlling", "difficult" and "not trying" can close down curiosity. Behaviour communicates something even when the person cannot yet explain it. Understanding is not the same as approving every behaviour; it means responding to the need and the consequences more intelligently. A particularly important point here is that understanding what the behaviour changes in the moment can reveal what support is actually needed; self-harm and suicide are not identical although risk must always be considered. That changes the way we respond. Instead of arguing with the person or demanding immediate change, we can look at what keeps the cycle going and what would create enough safety or flexibility for a different response to become possible. People also deserve to be believed about the effort involved. A task that looks small from the outside may involve preparation, sensory management, intrusive thoughts, fear, shame or recovery afterwards. Support becomes more effective when it is based on actual capacity rather than assumptions about what someone "should" manage. It can also help to notice what happens when the person is given time, choice and a sense of safety. With self-harm, pressure can make a response look more fixed, while calm curiosity may reveal needs that were hidden by fear or overload. This is why support should be judged by whether it increases safety, understanding and sustainable choice, not simply by whether it produces immediate compliance. There is rarely one perfect strategy. What helps on one day may be too much on another, and progress can include setbacks without meaning that everything has been lost. Keeping the person involved in decisions, reviewing what actually helped, and adapting support when circumstances change are all part of respectful care.

---

## What this can do to everyday life

Self-Harm can affect far more than the symptom that first attracts attention. It can change concentration, memory, decision-making, sleep, appetite, energy, confidence and relationships. It may alter how someone travels, shops, works, eats, socialises, attends healthcare or copes with unexpected change. Over time, life can quietly become organised around avoiding distress. Relationships can become caught in the pattern too. Loved ones may reassure, take over tasks, monitor, argue, withdraw or become frightened. These responses are understandable, especially when everyone is tired, but they can sometimes maintain the cycle or increase shame. Families and partners often need information and support in their own right. Work and education can be affected even when attendance remains good. Concentration may take more effort, meetings can use up capacity, deadlines may trigger spirals, and recovery may happen entirely outside working hours. Reasonable adjustments, predictable communication, quieter spaces, flexibility and clearer expectations can reduce unnecessary load. There can also be a secondary layer of distress created by misunderstanding the original problem. People may start believing they are weak, unreliable, selfish or failing. Separating the person from the pattern is important: a symptom or coping response is something a person is experiencing or doing, not their entire identity. Tracking patterns can help when it is done gently rather than obsessively. Useful things to notice include what happened beforehand, body sensations, thoughts or images, urges, what the person did next, the short-term effect and the longer-term cost. This can reveal the point in the cycle where support is most useful. It can also help to notice what happens when the person is given time, choice and a sense of safety. With self-harm, pressure can make a response look more fixed, while calm curiosity may reveal needs that were hidden by fear or overload. This is why support should be judged by whether it increases safety, understanding and sustainable choice, not simply by whether it produces immediate compliance. There is rarely one perfect strategy. What helps on one day may be too much on another, and progress can include setbacks without meaning that everything has been lost. Keeping the person involved in decisions, reviewing what actually helped, and adapting support when circumstances change are all part of respectful care.

---

## What we can learn from understanding this

The first lesson is that behaviour makes more sense when we understand its function. With self-harm, the visible response may be an attempt to find safety, certainty, relief, control, numbness, escape or connection. If we respond only to the surface behaviour, we can miss the problem the person is actually trying to solve. The second lesson is that understanding what the behaviour changes in the moment can reveal what support is actually needed; self-harm and suicide are not identical although risk must always be considered. This invites a different kind of compassion: not pity, and not pretending everything is fine, but an accurate understanding of why change can be difficult. Compassion and accountability can exist together. We can acknowledge impact while still asking what support makes a safer response more achievable. The third lesson is that early patterns matter. When we learn to recognise increasing avoidance, shrinking routines, rising reassurance, loss of recovery time, worsening sleep, increasing secrecy or a growing sense of hopelessness, support can begin before a crisis. Earlier support is often less about having the perfect words and more about noticing change and staying curious. The fourth lesson is that environment matters. We often ask people to regulate themselves inside situations that remain overwhelming, inaccessible or unsafe. Sometimes the most therapeutic change is practical: fewer demands, clearer communication, food that feels safe, predictable plans, financial advice, workplace adjustments, sensory changes or help getting to an appointment. Finally, recovery is not measured only by the disappearance of symptoms. It can also mean more choice, less secrecy, greater flexibility, better understanding of triggers, stronger boundaries, improved relationships, asking for help earlier and having a life that is no longer organised entirely around fear or coping. It can also help to notice what happens when the person is given time, choice and a sense of safety. With self-harm, pressure can make a response look more fixed, while calm curiosity may reveal needs that were hidden by fear or overload. This is why support should be judged by whether it increases safety, understanding and sustainable choice, not simply by whether it produces immediate compliance. There is rarely one perfect strategy. What helps on one day may be too much on another, and progress can include setbacks without meaning that everything has been lost. Keeping the person involved in decisions, reviewing what actually helped, and adapting support when circumstances change are all part of respectful care.

## What self-harm can teach us

Self-harm teaches us to look beneath behaviour rather than stopping at the shock of seeing it. The act may be serving a function: reducing unbearable arousal, interrupting numbness, expressing distress, creating a sense of control or responding to self-directed anger. Understanding the function does not make self-harm safe; it gives us better information for finding alternatives that meet the underlying need.

## Connection without surveillance

Support can become so focused on preventing another episode that the person feels watched rather than understood. Safety matters, but so do dignity and trust. Collaborative safety planning, direct questions about suicide when needed, appropriate wound or medical care and calm follow-up are more useful than punishment or threats. Families also need somewhere to process their fear so that the person who self-harms is not required to manage everybody else's distress.

---

## Support that is more likely to help

Support should be collaborative. Ask what the person finds difficult, what they already know about themselves and what kind of help feels possible. Person-centred psychosocial assessment, collaborative safety planning and tailored psychological support are important; urgent medical or crisis help may sometimes be needed. The best option depends on severity, history, physical health, previous treatment, access needs and the person's preferences. For supporters, listening before fixing is often a good starting point. Avoid debates designed to prove the person wrong while they are highly distressed. Keep communication clear and calm. Ask whether they want company, practical help, help making a plan, or simply somebody to stay present. If a strategy is part of a treatment plan, try to support that plan consistently rather than creating a new response every time distress rises. Support should also be accessible. Neurodivergent people may need adjustments for sensory differences, communication, interoception, executive functioning or masking. People with hearing, literacy or language needs may need information in another format. A treatment is not truly available if the person cannot access or process the way it is delivered. Self-support can begin with reducing shame and increasing observation. Notice patterns without turning the exercise into another test to pass. Protect basic needs where possible: sleep, hydration, food, medication, movement, rest and contact with safe people. Small sustainable changes are usually more useful than a burst of effort followed by collapse. If symptoms are persistent, worsening, substantially restricting life or becoming unsafe, professional assessment is appropriate. A GP can consider physical and mental health together and discuss treatment or referral. Some common mental-health difficulties can be self-referred to NHS Talking Therapies in England, while other conditions require specialist pathways. It can also help to notice what happens when the person is given time, choice and a sense of safety. With self-harm, pressure can make a response look more fixed, while calm curiosity may reveal needs that were hidden by fear or overload. This is why support should be judged by whether it increases safety, understanding and sustainable choice, not simply by whether it produces immediate compliance. There is rarely one perfect strategy. What helps on one day may be too much on another, and progress can include setbacks without meaning that everything has been lost. Keeping the person involved in decisions, reviewing what actually helped, and adapting support when circumstances change are all part of respectful care.

---

## When things feel more serious

It is important not to use a downloadable guide to diagnose yourself or somebody else. Self-Harm can overlap with other mental and physical health conditions, and new or severe physical symptoms deserve appropriate medical assessment. Professional help is particularly important when daily functioning is falling, distress is escalating, eating or sleeping is significantly affected, substances are being used to cope, or the person is becoming increasingly isolated. Ask directly about safety when you are worried. Talking about suicidal thoughts does not plant the idea. If somebody says they may act on suicidal thoughts, has a serious injury, has taken an overdose, or cannot keep themselves safe, urgent help is needed. In England, NHS 111 can provide urgent mental-health help. Call 999 or attend A&E; in an immediate life-threatening emergency. Samaritans can be contacted free on 116 123 for confidential emotional support. For self-harm or eating difficulties, physical risk may need medical attention even when the person does not describe themselves as suicidal or "ill enough". For trauma, ongoing danger must be addressed rather than treating every response as though the threat exists only in memory. For panic or health anxiety, appropriate medical assessment and anxiety treatment are not mutually exclusive. The message underneath this whole series is simple: understanding changes support. When we know more about what a response is doing, we can replace judgement with better questions, recognise risk earlier and help people find options that fit their real lives. Further support: NHS mental-health information and local services; NICE guidance for evidence-based treatment; Mind for mental-health information and support; Samaritans 116 123 for emotional support. This resource is educational and is not a substitute for individual medical or psychological care. It can also help to notice what happens when the person is given time, choice and a sense of safety. With self-harm, pressure can make a response look more fixed, while calm curiosity may reveal needs that were hidden by fear or overload. This is why support should be judged by whether it increases safety, understanding and sustainable choice, not simply by whether it produces immediate compliance. There is rarely one perfect strategy. What helps on one day may be too much on another, and progress can include setbacks without meaning that everything has been lost. Keeping the person involved in decisions, reviewing what actually helped, and adapting support when circumstances change are all part of respectful care.