

BRIGHT FUTURES HANDOUT ► PARENT

6 MONTH VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.

✓ HOW YOUR FAMILY IS DOING

- If you are worried about your living or food situation, talk with us. Community agencies and programs such as WIC and SNAP can also provide information and assistance.
- Don't smoke or use e-cigarettes. Keep your home and car smoke-free. Tobacco-free spaces keep children healthy.
- Don't use alcohol or drugs.
- Choose a mature, trained, and responsible babysitter or caregiver.
- Ask us questions about child care programs.
- Talk with us or call for help if you feel sad or very tired for more than a few days.
- Spend time with family and friends.

✓ YOUR BABY'S DEVELOPMENT

- Place your baby so she is sitting up and can look around.
- Talk with your baby by copying the sounds she makes.
- Look at and read books together.
- Play games such as peekaboo, patty-cake, and so big.
- Don't have a TV on in the background or use a TV or other digital media to calm your baby.
- If your baby is fussy, give her safe toys to hold and put into her mouth. Make sure she is getting regular naps and playtimes.

✓ FEEDING YOUR BABY

- Know that your baby's growth will slow down.
- Be proud of yourself if you are still breastfeeding. Continue as long as you and your baby want.
- Use an iron-fortified formula if you are formula feeding.
- Begin to feed your baby solid food when he is ready.
- Look for signs your baby is ready for solids. He will
 - Open his mouth for the spoon.
 - Sit with support.
 - Show good head and neck control.
 - Be interested in foods you eat.

Starting New Foods

- Introduce one new food at a time.
- Use foods with good sources of iron and zinc, such as
 - Iron- and zinc-fortified cereal
 - Pureed red meat, such as beef or lamb
- Introduce fruits and vegetables after your baby eats iron- and zinc-fortified cereal or pureed meat well.
- Offer solid food 2 to 3 times per day; let him decide how much to eat.
- Avoid raw honey or large chunks of food that could cause choking.
- Consider introducing all other foods, including eggs and peanut butter, because research shows they may actually prevent individual food allergies.
- To prevent choking, give your baby only very soft, small bites of finger foods.
- Wash fruits and vegetables before serving.
- Introduce your baby to a cup with water, breast milk, or formula.
- Avoid feeding your baby too much; follow baby's signs of fullness, such as
 - Leaning back
 - Turning away
- Don't force your baby to eat or finish foods.
 - It may take 10 to 15 times of offering your baby a type of food to try before he likes it.

Helpful Resources: Smoking Quit Line: 800-784-8669 | Poison Help Line: 800-222-1222

Information About Car Safety Seats: www.safercar.gov/parents | Toll-free Auto Safety Hotline: 888-327-4236

6 MONTH VISIT—PARENT



HEALTHY TEETH

- Ask us about the need for fluoride.
- Clean gums and teeth (as soon as you see the first tooth) 2 times per day with a soft cloth or soft toothbrush and a small smear of fluoride toothpaste (no more than a grain of rice).
- Don't give your baby a bottle in the crib. Never prop the bottle.
- Don't use foods or juices that your baby sucks out of a pouch.
- Don't share spoons or clean the pacifier in your mouth.

WHAT TO EXPECT AT YOUR BABY'S 9 MONTH VISIT

We will talk about

- Caring for your baby, your family, and yourself
- Teaching and playing with your baby
- Disciplining your baby
- Introducing new foods and establishing a routine
- Keeping your baby safe at home and in the car



SAFETY

- Use a rear-facing-only car safety seat in the back seat of all vehicles.
- Never put your baby in the front seat of a vehicle that has a passenger airbag.
- If your baby has reached the maximum height/weight allowed with your rear-facing-only car seat, you can use an approved convertible or 3-in-1 seat in the rear-facing position.
- Put your baby to sleep on her back.
- Choose crib with slats no more than 2 3/8 inches apart.
 - Lower the crib mattress all the way.
- Don't use a drop-side crib.
- Don't put soft objects and loose bedding such as blankets, pillows, bumper pads, and toys in the crib.
- If you choose to use a mesh playpen, get one made after February 28, 2013.
- Do a home safety check (stair gates, barriers around space heaters, and covered electrical outlets).
- Don't leave your baby alone in the tub, near water, or in high places such as changing tables, beds, and sofas.
- Keep poisons, medicines, and cleaning supplies locked and out of your baby's sight and reach.
- Put the Poison Help line number into all phones, including cell phones. Call us if you are worried your baby has swallowed something harmful.
- Keep your baby in a high chair or playpen while you are in the kitchen.
- Do not use a baby walker.
- Keep small objects, cords, and latex balloons away from your baby.
- Keep your baby out of the sun. When you do go out, put a hat on your baby and apply sunscreen with SPF of 15 or higher on her exposed skin.

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th Edition

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.

Suggested Feeding Schedule

Age	Breast Milk or Formula	Grain Products	Juices	Vegetables	Fruits	Protein-Rich Foods
Birth – 4 Month	Breast: 8-12+ feedings	None	None	None	None	None
4-6 Months	Iron-Fortified Infant Formula: 14-42 ounces Breast: 5 or more feedings	None	None	None	None	None
6-8 Months	Iron-Fortified Infant Formula: 26-39 ounces Breast: 3-5 feedings	Iron fortified infant cereals or enriched hot cereals (4-6Tbsp)	None	Plain strained or pureed fresh or cooked vegetables (3-4 Tbsp)	Plain strained or pureed fresh or cooked fruit (3-4 Tbsp)	Plain strained or pureed protein-rich foods such as meats, egg, fish, cheese, yogurt and peanut butter (1-3 tbsp)
	Iron-Fortified Infant Formula: 24-32 ounces	Dry toast, small pieces of crackers, or dry breakfast cereals and other grain products (4-6 Tbsp)				(If it's not a choking hazard)
8-12 Months	Breast: 3-4 feedings	Iron-fortified infant cereals or enriched hot cereals (4-6 Tbsp)	None	Plain strained or pureed fresh or cooked vegetables (3-4 Tbsp)	Plain strained or pureed fresh or cooked fruit (3-4 Tbsp)	Pureed or chopped protein-rich foods such as meats, egg, fish, cheese, yogurt and peanut butter (1-3 tbsp)
	Iron-Fortified Infant Formula: 24-32 ounces	Dry toast, small pieces of crackers, or dry breakfast cereals and other grain products (4-6 Tbsp)				(If it's not a choking hazard)
Comments	By about 11 to 12 months, try to wean entirely off the bottle and onto a cup.	Never add honey to cereal or any foods Avoid foods that may cause choking	Avoid feeding soda, juices, fruit punches, ades and drinks, gelatin, water, coffee or tea.	It is not necessary to add salt, sugar, oil, butter, other fats or seasoning Avoid foods that may cause choking	Do not add sugar or syrups to fruits Never add honey to fruit or any foods Remove seeds and pits from fruits	Avoid fried meats, gravies, sauces, processed meats Check carefully for bones



Where We Stand: Vitamin D & Iron Supplements for Babies

The American Academy of Pediatrics (AAP) recommends infants be fed breast milk (also known as human milk) exclusively for the first 6 months after birth. Human milk contains a natural balance of vitamins, especially C, E and the B vitamins. So, if you and your baby are both healthy, and you are well nourished, your child may not require any supplements of these vitamins. However, breastfed infants need supplemental vitamin D.



Why do babies need vitamin D supplements?

Our bodies need sunlight to produce vitamin D, which is why exposure to sunlight is good in moderation. However, children should wear sunscreen, hats, and protective clothing when outdoors for extended periods of time to prevent sunburn and reduce the risk of skin cancer later in life. For this reason, the AAP recommends that all infants receive a vitamin D supplement (unless they are consuming more than 27 ounces per day of commercial formula that has the vitamin D supplement added).

How much vitamin D do babies need?

Vitamin D supplements of 400 IU (10 mcg) per day are recommended for babies up until age one year, with 600 IU (15 mcg) per day for children over one year. Talk to your pediatrician about supplemental vitamin D drops.

Do babies need iron supplements?

For the first four months, your breastfed baby needs no additional iron. The iron in their body at birth was enough for their initial growth. But now the reserves will be low and as their growth increases, so will their need for iron. At four months of age infants who are partially or completely breastfeeding should be supplemented with 1 mg/kg per day of oral iron until appropriate iron-containing complementary foods (including iron-fortified cereals) are introduced in their diet.

Universal screening for iron deficiency

The AAP recommends that all babies be screened at 12 months of age for iron deficiency and iron deficiency anemia.

See [Anemia and Your Child: Parent FAQs \(/English/health-issues/conditions/chronic/Pages/Anemia-and-Your-Child.aspx\)](/English/health-issues/conditions/chronic/Pages/Anemia-and-Your-Child.aspx) for more information.

If there were pregnancy or birth complications such as diabetes, low birth weight, or prematurity, or if your baby was small for gestational age and is taking breast milk, iron supplementation may start in the first month after birth. Fortunately, once you start your baby on solid foods, they'll also receive iron from meats, iron-fortified cereals and green vegetables.

More information

- [Vitamin D Deficiency and Rickets \(/English/healthy-living/nutrition/Pages/Vitamin-D-Deficiency-and-Rickets.aspx\)](/English/healthy-living/nutrition/Pages/Vitamin-D-Deficiency-and-Rickets.aspx)
- [Where We Stand: Vitamins \(/English/healthy-living/nutrition/Pages/Where-We-Stand-Vitamins.aspx\)](/English/healthy-living/nutrition/Pages/Where-We-Stand-Vitamins.aspx)

[ps to Reduce Arsenic in Your Baby's Diet \(/English/ages-stages/baby/feeding-nutrition/Pages/reduce-arsenic.aspx\)](/English/ages-stages/baby/feeding-nutrition/Pages/reduce-arsenic.aspx)

[Back to Top](#)

- [Dietary Supplements for Toddlers \(/English/ages-stages/toddler/nutrition/Pages/Dietary-Supplements-for-Toddlers.aspx\)](#)

Last Updated 5/24/2022

Source Adapted from Caring for Your Baby and Young Child: Birth to Age 5 7th Edition (Copyright © 2019 American Academy of Pediatrics)

The information contained on this Web site should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

Sleeping Well: Tips for Parents of Babies and Young Children

It's important for children to develop good sleep habits early on. Sleeping well is important to children's development and well-being. Here is information from the American Academy of Pediatrics to help ensure a good night's sleep for children—and parents too! Remember that children differ in how much sleep they need, how long it takes them to fall asleep, and how easily they wake up. If you have any questions about your child's sleep habits, ask your child's doctor.

Babies

Babies do not have regular sleep cycles until about 4 months of age. Although newborns sleep about 16 to 17 hours per day, they may sleep for only 1 or 2 hours at a time. As babies get older, they need less sleep. Keep in mind that your baby's sleep needs and patterns may differ from those of other babies.

How to Help Babies Sleep Better at Night

All Babies

- Try not to stimulate or wake babies when you feed or change them during the night. If you speak, speak softly.
- Make daytime playtime. Keep babies awake longer during the day. This helps babies sleep for longer periods during the night. Spend time talking, reading, and playing together.

Babies 4 Months and Older

- Put babies to bed when they are drowsy. Do not wait until babies are asleep. This helps babies learn to fall asleep on their own, in their own bed. If you hold babies or rock them to sleep, they may struggle to go back to sleep if they wake up during the night. Remember to place babies on their back to sleep for every sleep until they are 1 year of age. Visit www.HealthyChildren.org/safesleep to learn more about how to create a safe sleeping environment for babies.
- Do not rush in to soothe a crying baby. Babies need time to put themselves back to sleep, and they need to learn how to fall back asleep on their own. It is normal for a 6-month-old to wake up during the night and then go back to sleep after a few minutes. Of course, you can attend to them—like feeding them, changing a soiled diaper, or comforting them if they are sick—if needed.

Toddlers and Preschoolers

Many parents find their toddler's bedtime to be the hardest part of the day. Children this age often resist going to sleep, especially if they have older siblings who are still awake.

How to Help Toddlers Develop Good Sleep Habits

- Set up a quiet routine before bedtime. This helps children understand that it will soon be time to go to sleep. Use this time to read a story, put on quiet music, or give a bath. Do not play with children before bed, because active play may make them too excited to sleep.

- Be consistent. Make bedtime the same time every night. This helps children know what to expect and helps them establish healthy sleep patterns.
- Let children take a favorite thing to bed each night. Children may want to sleep with a teddy bear, special blanket, or favorite toy. This often helps children fall asleep, especially if they wake up during the night. Make sure the object is safe and doesn't have any choking hazards, like buttons or loose ribbons. Stuffing or pellets inside stuffed toys can also be dangerous.
- Make sure children are comfortable. Take care of children's needs before bedtime so they don't use them as reasons to avoid going to sleep. Children may want a drink of water, a light left on, or the door left slightly open.
- Do not let children sleep in the same bed with you. This can make it harder for them to fall asleep when they are alone.
- Do not return to a child's room when they call out. Instead, try to
 - Make sure children are safe and well before bedtime. If they are safe and well, there is no need to go into their room while they sort out how to get themselves back to sleep.
 - Keep in mind that children's main goal is to get you to appear. So if you appear for any reason, even just for "checking," they will expect you to come each time they call out.
- Give it time. Helping children develop good sleep habits can be a challenge, and it is normal to get upset when children keep you awake at night. Try to be understanding. A negative response from a parent can sometimes make a sleep problem worse. Keep in mind that children need the time and opportunity to find out how go back to sleep on their own when they wake up during the night.

For More Information

American Academy of Pediatrics

www.aap.org and www.HealthyChildren.org

Adapted from the American Academy of Pediatrics (AAP) brochure *Sleep Problems in Children*.

The AAP is an organization of 67,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of all infants, children, adolescents, and young adults.

In all aspects of its publishing program (writing, review, and production), the AAP is committed to promoting principles of equity, diversity, and inclusion.

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



healthychildren.org

Powered by pediatricians. Trusted by parents.
from the American Academy of Pediatrics

Children's Dental Health: What You Need to Know

A bright smile begins long before the first tooth appears. Parent and caregiver help is important for children to develop healthy teeth. Read on for information from the American Academy of Pediatrics about caring for your child's teeth.

Steps to good dental health include

- Regular care by a dentist beginning by 1 year of age
- Enough fluoride (in water, toothpaste, and fluoride varnish)
- Brushing and flossing 2 times each day
- Eating healthy and limiting sugar
- Using a mouth guard during sports participation to prevent injury, if necessary

Fluoride is important because it

- Hardens tooth enamel (the outside coating on teeth)
- Repairs early damage to teeth all day every day

Note: Fluoride is a natural substance that can be added to drinking water, toothpaste, mouthwash, and varnish (dental treatment). During well-child visits (also known as health supervision visits), doctors may recommend drinking more fluoridated water or, for some children, using fluoride tablets or drops. Also, fluoride varnish should be applied to children's teeth by their doctor or dentist up to 4 times per year.

Here's how to clean your child's teeth.

Babies to 3 years of age

- Wipe the gums 2 times each day with a piece of gauze or a damp cloth until the first tooth or teeth arrive.
- Brush the first tooth or teeth with a soft toothbrush 2 times each day. Brush for 2 minutes each time.
- Use "Just a dot, not a lot!" of fluoride toothpaste. The amount of toothpaste should be the size of a small grain of rice for children younger than 3 years.

Children 3 years and older

- Brush your child's teeth with a soft toothbrush 2 times each day. Brush for 2 minutes each time. Children should learn how to brush their teeth on their own. However, parents should brush their children's teeth first before handing over the toothbrush to their children until they are 7 years of age.
- Use "Just a dot, not a lot!" of fluoride toothpaste. The amount of toothpaste should be the size of a small pea for children 3 years and older.

All children

- Teach your child to spit out excess toothpaste. Your child may want to swallow the toothpaste because it tastes good. However, swallowing too much toothpaste can result in white spotting of the teeth called fluorosis. Children should not rinse after brushing and spitting out excess toothpaste.

- Floss where any 2 teeth touch each other to prevent a cavity forming between the teeth.
- Check front and back of the teeth for early signs of tooth decay, such as white, yellow, or brown spots or lines on the teeth. Lift up the top lip to get a good look at the front upper teeth.
- Change your child's toothbrush every 6 months.

Here are other ways to help prevent tooth decay in babies and children.

- Schedule regular dental checkups for each family member.
- Avoid sharing food, drinks, spoons, and forks. If your baby is using a pacifier, avoid licking it to clean it.
- Offer water if your child is thirsty. Also, only offer water in sippy cups between meals and in bedtime bottles. Sipping juices, sports drinks, flavored drinks, lemonade, soft drinks (soda, pop), or flavored teas throughout the day causes acid attacks on teeth.
- Offer healthy snacks such as fruits or vegetables. Avoid offering sweet or sticky snacks, such as raisins, gummy candies, and vitamins, or fruit-flavored snacks/rolls or cookies. There is sugar in foods like crackers and chips too. Reserve these for desserts at the end of meals.
- Be sure to clean teeth after your child drinks milk at bedtime.
- Check front and back of the teeth for early signs of tooth decay—white, yellow, or brown spots or lines on the teeth. Lift your child's lip to get a better look at the upper front teeth. This should be done about once a month.

Common Questions

Does pacifier use or thumb-sucking hurt teeth?

Sucking on a pacifier, thumb, or fingers may affect the shape of the mouth or how teeth are lining up.

- If the habit stops by 3 years of age, the teeth will usually correct themselves without treatment.
- If the sucking habit continues after "permanent" teeth have come in, orthodontic care may be needed to line the teeth up for disease prevention and appearance.

What should I do when my child falls and loosens a tooth?

- Call your child's dentist or pediatrician for advice.
- For the next 6 months or so, watch for redness in the gum above the loosened tooth and notify your child's dentist immediately if it occurs.

What is a pediatric dentist?

Pediatric dentists have special training to provide routine dental care for children and can care for children with complicated oral health problems. They are specialists in the care of children's teeth and mouth problems, especially when

- Teeth are chipped or injured or there is an injury in the mouth area.
- Teeth show signs of discoloration that could be tooth decay or trauma.

- Children complain of tooth pain or sensitivity to hot or cold foods or liquids. This could also be a sign of decay.
- There is any abnormal growth inside the mouth.
- Children have an unusual bite (in other words, their teeth do not fit together right).

You can find a pediatric dentist in your area on the American Academy of Pediatric Dentistry website at www.aapd.org. Some family dentists are trained to provide care for children without complicated problems. There are also many family dentists who provide preventive care to healthy children.

When should my child begin regular dental checkups?

- All infants should receive oral health risk assessments by 6 months of age at their well-child visit with their medical provider and at every well-child checkup.
- Children with special health care needs should be referred to a dentist as early as 6 months of age, and no later than 12 months of age, to establish their dental home and may be seen more frequently than typical children.
- Every child should have a dental home established by 12 months of age.

Remember

If you have any questions or concerns about your child's teeth, contact your child's dentist.

American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN®



healthychildren.org
Powered by pediatricians. Trusted by parents.
from the American Academy of Pediatrics

The American Academy of Pediatrics (AAP) is an organization of 67,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

Any websites, brand names, products, or manufacturers are mentioned for informational and identification purposes only and do not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of external resources. Information was current at the time of publication. The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

© 2020 American Academy of Pediatrics. All rights reserved.



BIRTH TO 6 MONTHS

Safety for Your Child

Did you know that hundreds of children younger than 1 year die every year in the United States because of injuries—most of which could be prevented?

Often, injuries happen because parents are not aware of what their children can do. Children *learn quickly*, and before you know it, your child will be *wiggling* off a bed or *reaching* for your cup of hot coffee.

Car Injuries

Car crashes are a great threat to your child's life and health. Most injuries and deaths from car crashes **can be prevented** by the use of car safety seats. Your child, besides being much safer in a car safety seat, will be more comfortable and content, so you can pay attention to your driving. Make your newborn's first ride home from the hospital a safe one—in a car safety seat. Your infant should ride in the back seat in a rear-facing car safety seat.

Make certain that your baby's car safety seat is installed correctly. Read and follow the instructions that come with the car safety seat and the sections in the owners' manual of your car on using car safety seats correctly. Use the car safety seat EVERY time your child is in the car.

NEVER put an infant in the front seat of a car with a passenger airbag.



Falls

Babies *wiggle* and *move* and *push* against things with their feet soon after they are born. Even these very first movements can result in a fall. As your baby grows and is able to roll over, they may fall off of things unless protected. **Do not leave your baby alone** on changing tables, beds, sofas, or chairs. **Put your baby in a safe place** such as a crib or playpen when you cannot hold them.

Your baby may be able to crawl as early as 6 months. **Use gates on stairways and close doors** to keep your baby out of rooms where they might get hurt. **Install operable window guards** on all windows above the first floor.

Do not use a baby walker. Your baby may tip the walker over, fall out of it, or fall down stairs and seriously injure their head. Baby walkers let children get to places where they can pull heavy objects or hot food on themselves. A better choice is a stationary activity center with no wheels.

If your child has a serious fall or does not act normally after a fall, call your doctor.



(over)

Burns

At 3 to 5 months, babies will wave their fists and grab at things. **NEVER carry your baby and hot liquids, such as coffee, or foods at the same time.** You can't handle both. Your baby can get burned! To protect your child from tap water scalds, the hottest temperature at the faucet should be no more than 120 °F (48.9 °C). In many cases you can adjust your water heater.

If your baby gets burned, immediately put the burned area in cold water. Keep the burned area in cold water for a few minutes to cool it off. Then cover the burn loosely with a dry bandage or clean cloth and call your doctor.

To protect your baby from house fires, be sure you have a working smoke alarm on every level of your home, especially in furnace and sleeping areas. Test the alarms every month. It is best to use smoke alarms that use long-life batteries, but if you do not, change the batteries at least once a year.




Choking and Suffocation

Babies explore their environment by putting anything and everything into their mouths. **NEVER leave small objects in your baby's reach, even for a moment.** If you start transitioning to solid food, NEVER feed your baby hard pieces of food such as chunks of raw carrots, apples, hot dogs, grapes, peanuts, and popcorn. Cut all the foods you feed your baby into thin pieces to prevent choking. **Be prepared if your baby starts to choke. Ask your doctor to recommend the steps you need to know. Learn how to save the life of a choking child.**

To prevent possible suffocation and reduce the risk of sudden infant death syndrome (SIDS), **your baby should always sleep on their back. Your baby should have their own crib or bassinet with no pillows, stuffed toys, bumpers, or loose bedding. NEVER put your baby on a waterbed, beanbag, or anything that is soft enough to cover the face and block air to the nose and mouth.**

Plastic wrappers and bags form a tight seal if placed over the mouth and nose and may suffocate your child. Keep them away from your baby.




From Your Doctor

The American Academy of Pediatrics (AAP) is an organization of 67,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of all infants, children, adolescents, and young adults.

In all aspects of its publishing program (writing, review, and production), the AAP is committed to promoting principles of equity, diversity, and inclusion.

The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

Car Safety Seat Checkup



Using a car safety seat correctly makes a big difference. It is important to choose the right seat for your child's size and make sure it is used correctly to properly protect your child in a crash. Here are car safety seat tips from the American Academy of Pediatrics.

Does your car have airbags?

- Never place a rear-facing car safety seat in the front seat of a vehicle that has a front passenger airbag. If the airbag inflates, it will hit the back of the car safety seat, right where your baby's head rests, and could cause serious injury or death.
- The safest place for all children younger than 13 years to ride is in the back seat regardless of weight and height.
- If an older child must ride in the front seat, a child in a forward-facing car safety seat with a harness may be the best choice. Be sure you move the vehicle seat as far back from the dashboard (and airbag) as possible.

Is your child facing the right way for weight, height, and age?

- All infants and toddlers should ride in a rear-facing car safety seat for as long as possible, up to the highest weight or height allowed by their car safety seat manufacturer. Most infants will outgrow rear-facing-only seats by length long before they reach the maximum weight limit. When infants outgrow a rear-facing-only seat, they should use a rear-facing convertible seat. Most convertible seats have limits that will allow children to ride rear facing for 2 years or more.
- When a child exceeds the rear-facing weight or height limit for their convertible car safety seat, they should use a forward-facing seat with a harness for as long as possible, up to the highest weight or height allowed by their car safety seat manufacturer. Most seats can accommodate children up to 65 pounds or more.

Is the harness snug?

- Harness straps should fit snugly against your child's body. Check the car safety seat instructions to learn how to adjust the straps.
- Place the chest clip at armpit level to keep the harness straps secure on the shoulders.

Does the car safety seat fit correctly in your vehicle?

- Not all car safety seats fit properly in all vehicles.
- Read the section on car safety seats in the owner's manual for your car.

Should you use the LATCH system?

- Car safety seats may be installed with either the vehicle's seat belt or its LATCH (lower anchors and tethers for children) system. Both systems are equally safe, so caregivers should use one or the other, whichever works best for them, their car safety seat, and their vehicle. In general, caregivers should use only 1 of the 2 options, unless the car safety seat and vehicle manufacturers say it is OK to use both systems at the same time.
- Vehicles with the LATCH system have anchors located in the back seat, where the seat cushions meet. All car safety seats have attachments that fasten to these anchors. Nearly all passenger vehicles made on or after September 1, 2002, and all car safety seats are equipped to use LATCH. All lower anchors are rated for a maximum weight of 65 pounds (total weight includes car safety seat and child). Check the car safety seat manufacturer's recommendations for the maximum weight a child can be to use lower anchors. New car safety seats have the maximum weight printed on their label.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®





FIGURE 4.
Forward-facing car safety seat with
harness.



FIGURE 5.
Belt-positioning booster seat.



FIGURE 6.
Lap and shoulder seat belt.

Should you use a top tether?

- The top tether does a great job of improving safety provided by the seat. Use the tether for all forward-facing seats up to the highest weight allowed by your vehicle and the manufacturer of your car safety seat. Your vehicle owner's manual will tell you how to find the location of tether anchors. Remember, weight limits are different for different car safety seats and different vehicles, so it is important to check both the car safety seat and vehicle manufacturer instructions for lower anchors and tethers.

Is the seat belt or LATCH strap in the right place and pulled tight?

- Route the seat belt or LATCH strap through the correct belt path for the car safety seat. Convertible seats have different belt paths for when they are used rear facing or forward facing (check your instructions to make sure).
- If you install the car safety seat by using your vehicle's seat belt, you must make sure the seat belt locks to keep a tight fit. In most newer cars, you can lock the seat belt by pulling it all the way out and then allowing it to retract to keep the seat belt tight around the car safety seat. Many car safety seats have built-in lock-offs to lock the belt. Check your vehicle owner's manual and car safety seat instructions to make sure you are using the seat belt correctly.
- Pull the belt tight. Apply weight into the seat with your hand while tightening the seat belt or LATCH strap. When the car safety seat is installed, be sure it does not move more than an inch side to side or toward the front of the car.

Has your child outgrown the forward-facing seat?

- All children whose weight or height is above the forward-facing limit for their car safety seat should use a belt-positioning booster seat until the vehicle seat belt fits properly. While this is typically when they have reached at least 4 feet 9 inches in height and are 8 to 12 years of age,

every kid is different. It is important to make sure to check how a seat belt fits your child in each car they will be riding in.

- A seat belt fits properly when the shoulder belt lies across the middle of the chest and shoulder, not the neck or throat; the lap belt is low and snug across the upper thighs, not the belly; and the child is tall enough to sit against the vehicle seat back with their knees bent over the edge of the seat without slouching and can comfortably stay in this position throughout the trip.

Has the car safety seat been recalled?

- You can find out by calling the manufacturer or the National Highway Traffic Safety Administration (NHTSA) Vehicle Safety Hotline at 888-327-4236 or by going to the NHTSA website at NHTSA.gov.
- Follow the manufacturer's instructions for making any repairs to your car safety seat.
- Be sure to fill in and mail in the registration card that comes with the car safety seat. It will be important in case the seat is recalled.

Do you know the history of your child's car safety seat?

- Do not use a used car safety seat if you do not know the history of the seat.
- Do not use a car safety seat that has been in a crash, has been recalled, is too old (check the expiration date or use 6 years from the manufacture date if there is no expiration date), has any cracks in its frame, or is missing parts.
- Make sure it has labels from the manufacturer and instructions. Keep the instructions with the car safety seat.
- Call the car safety seat manufacturer if you have questions about the safety of your seat.

Resources

If you have questions or need help installing your car safety seat, find a certified child passenger safety technician (CPST) by going to the National Child Passenger Safety Certification website at <http://cert.safekids.org> and clicking on "Find a Tech."

The American Academy of Pediatrics (AAP) offers more information in the brochure *Car Safety Seats: A Guide for Families*. Ask your pediatrician about this brochure or visit the official AAP website for parents, www.HealthyChildren.org/carseatguide.

Figure 1 adapted from US Department of Transportation, National Highway Traffic Safety Administration. *LATCH Makes Child Safety Seat Installation as Easy as 1-2-3*. DOT HS publication 809 489. March 2011. Figures 2, 3, 4, 5, and 6 by Anthony Alex LeTourneau.

Any websites, brand names, products, or manufacturers are mentioned for informational and identification purposes only and do not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of external resources. Information was current at the time of publication. The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

© 2024 American Academy of Pediatrics.
All rights reserved.



Choking Prevention and First Aid for Infants and Children

When children begin crawling or eating table foods, parents must be aware of the dangers and risks of choking. Children younger than 5 years can easily choke on food and small objects.

Choking occurs when food or small objects get caught in the throat and block the airway. This can prevent oxygen from getting to the lungs and the brain. When the brain goes without oxygen for more than 4 minutes, brain damage or even death may occur.

Many children die from choking each year, and some children who survive a severe choking episode have permanent, life-changing brain injuries. Most children who choke to death are younger than 5 years. Two-thirds of choking victims are infants younger than 1 year. Balloons, balls, marbles, pieces of toys, and foods cause the most choking deaths.

Read more about choking prevention and first aid.

Dangerous Foods

Keep dangerous foods from children until 4 years of age or older, depending on each child's development and maturity level. However, round, firm foods, such as hot dogs or grapes, can be served if completely chopped into tiny pieces. When infants and young children do not grind or chew their food well, they may try to swallow it whole. Peanut butter and other nut butters should be spread thinly.

Here are foods that can be choking hazards:

- Hot dogs
- Hard, gooey, or sticky candy
- Chewing gum
- Nuts and seeds
- Whole grapes
- Raw vegetables, such as carrot sticks
- Raw fruit chunks, such as apple chunks
- Popcorn
- Chunks of peanut butter or other nut butters
- Marshmallows
- Meat sticks/sausages
- Chunks of meat
- Chunks of cheese or string cheese

Dangerous Household Items

Keep the following household items away from infants and children:

- Balloons
- Coins
- Marbles

- Toys with small parts
- Toys that can be squeezed to fit entirely into a child's mouth
- Small balls
- Pen or marker caps
- Small button-type batteries
- Medicine syringes

What You Can Do To Prevent Choking

- Learn CPR (cardiopulmonary resuscitation) (basic life support).
- Be aware that balloons pose a choking risk to children up to 8 years of age.
- Keep dangerous foods from children until 4 years of age or older, depending on each child's development and maturity level.
- Insist that children eat at the table or sit down when they eat. They should never run, walk, play, or lie down with food in their mouths.
- Cut food for infants and young children into pieces no larger than one-half inch, and teach them to chew their food well.
- Supervise mealtime for infants and young children.
- Be aware of older children's actions. Many choking incidents occur when older brothers or sisters give dangerous foods, toys, or small objects to a younger child.
- Avoid toys with small parts, and keep other small household items out of the reach of infants and young children.
- Follow the age recommendations on toy packages. Age guidelines reflect the safety of a toy, which is based on any possible choking hazard, as well as the child's physical and mental abilities at various ages.
- Check under furniture and between cushions for small items that children could find and put in their mouths.
- Do not let infants and young children play with coins.

First Aid for the Child Who Is Choking

Make it a point to learn the instructions on the following pages of this publication. Post the chart in your home. However, these instructions should not take the place of an approved class in basic first aid, CPR, or emergency prevention. Contact your local American Red Cross (www.redcross.org) or the American Heart Association (www.heart.org) to find out about classes offered in your area. Most of the classes teach basic first aid, CPR, and emergency prevention, along with what to do for a choking infant or child. Your child's doctor also can help you understand these steps and talk with you about the importance of supervising mealtime and identifying dangerous foods and objects.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



healthychildren.org

Powered by pediatricians. Trusted by parents.
From the American Academy of Pediatrics

The American Academy of Pediatrics (AAP) is an organization of 66,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

The persons whose photographs are depicted in this publication are professional models. They have no relation to the issues discussed. Any characters they are portraying are fictional. Listing of resources does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of external resources. Information was current at the time of publication. The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

© 2018 American Academy of Pediatrics. All rights reserved.

CHOKING/CPR

LEARN AND PRACTICE CPR (CARDIOPULMONARY RESUSCITATION).

IF ALONE WITH A CHILD WHO IS CHOKING...

1. SHOUT FOR HELP. 2. START RESCUE EFFORTS. 3. CALL 911 OR YOUR LOCAL EMERGENCY NUMBER.

START FIRST AID FOR CHOKING IF

- The child cannot breathe at all (the chest is not moving up and down).
- The child cannot cough or talk or looks blue.
- The child is found unconscious/unresponsive. (Go to CPR.)

DO NOT START FIRST AID FOR CHOKING IF

- The child can breathe, cry, or talk.
- The child can cough, sputter, or move air at all. The child's normal reflexes are working to clear the airway.

FOR CHILDREN 1 TO 8 YEARS OF AGE

CHILD CHOKING (HEIMLICH MANEUVER)

Have someone call 911. If the child is choking and is unable to breathe, cough, cry, or speak, follow these steps.

1. Perform Heimlich maneuver.

- Place hand, made into a fist, and cover with other hand just above the navel. Place well below the bottom tip of the breastbone and rib cage.
- Give each thrust with enough force to produce an artificial cough designed to relieve airway obstruction.
- Perform Heimlich maneuver until the object is expelled or the child becomes unconscious/unresponsive.

2. If the child becomes UNCONSCIOUS/UNRESPONSIVE, begin CPR. →



CHILD CPR

To be used when the child is UNCONSCIOUS/UNRESPONSIVE or when breathing stops.
Place child on flat, hard surface.

1 START CHEST COMPRESSIONS.

- Place the heel of 1 or 2 hands over the lower half of the sternum.
- Compress chest at least $\frac{1}{3}$ the depth of the chest, or about 5 cm (2 inches).
- After each compression, allow chest to return to normal position. Compress chest at a rate of at least 100 to 120 times per minute.
- Do 30 compressions.



1-hand technique



2-hand technique


2 OPEN AIRWAY.

- Open airway (head tilt-chin lift).
- If you see a foreign body, sweep it out with your finger. Do NOT do blind finger sweeps.



3 START RESCUE BREATHING.

- Take a normal breath.
- Pinch the child's nose closed, and cover child's mouth with your mouth.
- Give 2 breaths, each for 1 second. Each breath should make the chest rise.



4 RESUME CHEST COMPRESSIONS.

- Continue with cycles of 30 compressions to 2 breaths until the object is expelled.
- After 5 cycles of compressions and breaths (about 2 minutes), if no one has called 911 or your local emergency number, call it yourself.

If at any time an object is coughed up or the infant/child starts to breathe, stop rescue breaths and call 911 or your local emergency number.

Ask your pediatrician for information on choking/CPR instructions for children older than 8 years and for information on an approved first aid or CPR course in your community.

CHOKING/CPR

LEARN AND PRACTICE CPR (CARDIOPULMONARY RESUSCITATION).

IF ALONE WITH A CHILD WHO IS CHOKING...

1. SHOUT FOR HELP. 2. START RESCUE EFFORTS. 3. CALL 911 OR YOUR LOCAL EMERGENCY NUMBER.

START FIRST AID FOR CHOKING IF

- The child cannot breathe at all (the chest is not moving up and down).
- The child cannot cough or talk or looks blue.
- The child is found unconscious/unresponsive. (Go to CPR.)

DO NOT START FIRST AID FOR CHOKING IF

- The child can breathe, cry, or talk.
- The child can cough, sputter, or move air at all. The child's normal reflexes are working to clear the airway.

FOR INFANTS YOUNGER THAN 1 YEAR

INFANT CHOKING

If the infant is choking and is unable to breathe, cough, cry, or speak, follow these steps. Have someone call 911.



ALTERNATING WITH




Alternate back blows (slaps) and chest compressions until the object is dislodged or the infant becomes unconscious/unresponsive. If the infant becomes unconscious/unresponsive, begin CPR.

INFANT CPR

To be used when the infant is **UNCONSCIOUS/UNRESPONSIVE** or when breathing stops. Place infant on flat, hard surface.


1 START CHEST COMPRESSIONS.

- Place 2 fingers of 1 hand on the breastbone just below the nipple line.
- Compress chest at least $\frac{1}{3}$ the depth of the chest, or about 4 cm (1.5 inches).
- After each compression, allow chest to return to normal position. Compress chest at rate of at least 100 times per minute.
- Do 30 compressions.




2 OPEN AIRWAY.

- Open the airway (head tilt-chin lift).
- If you see a foreign body, sweep it out with your finger. Do NOT do blind finger sweeps.




3 START RESCUE BREATHING.

- Take a normal breath.
- Cover infant's mouth and nose with your mouth.
- Give 2 breaths, each for 1 second. Each breath should make the chest rise.




4 RESUME CHEST COMPRESSIONS.

- Continue with cycles of 30 compressions to 2 breaths.
- After 5 cycles of compressions and breaths (about 2 minutes), and if no one has called 911 or your local emergency number, call it yourself.



If at any time an object is coughed up or the infant/child starts to breathe, stop rescue breaths and call 911 or your local emergency number.

Ask your pediatrician for information on choking/CPR instructions for children older than 8 years and for information on an approved first aid or CPR course in your community.



Pool Safety for Children

A swimming pool can be very dangerous for children. If possible, do not put a swimming pool in your yard while you have young children. Help protect your children from drowning by doing the following:

- Never leave your children alone in or near the pool, even for a moment. An adult who knows CPR should actively supervise children at all times.
- Practice touch supervision with children younger than 5 years and with children of any age who are not strong swimmers. This means that the adult is within an arm's length of the child at all times.
- You must put up a fence to separate your house from the pool. Most young children who drown in pools wander out of the house and fall into the pool when they are not expected to be swimming. Install a fence at least 4 feet high around all 4 sides of the pool. This fence must completely separate the pool from the house and play area of the yard. Use gates that self-close and self-latch, with latches higher than your children's reach.
- Keep rescue equipment (such as a shepherd's hook or life preserver) and a telephone by the pool.
- Do not use air-filled "swimming aids" as a substitute for approved life jackets.
- Remove all toys from the pool after use so children aren't tempted to reach for them.
- After the children are done swimming, secure the pool so they can't get back into it.
- Pool alarms and rigid safety covers that meet the standards of ASTM International may add to the protection of your children but should not be used in place of the fence between your house and the pool. Even fencing around your pool and using a rigid safety cover will not prevent all drownings.

While swim skills are an additional layer of prevention, remember that teaching your child how to swim DOES NOT mean your child is safe in water. Additional safety measures are necessary when your child is near the pool and when it is not swim time.

From Your Doctor

The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®





Home Water Hazards for Young Children

Each year, many young children drown in swimming pools, other bodies of water, and standing water around the home, such as

- Bathtubs
- Buckets and pails, especially 5-gallon buckets and diaper pails
- Ice chests with melted ice
- Toilets
- Hot tubs, spas, and whirlpools
- Irrigation ditches, postholes, and wells
- Fishponds and fountains

Children must be watched closely by an adult at all times when in or near water. Children may drown in an inch or 2 of water. Stay within an arm's length of your child.

Other safety activities include

- Empty all buckets, pails, and bathtubs completely after each use—do not leave them filled and unattended.
- Keep young children out of the bathroom unless they are closely watched. Teach others in the home to keep the bathroom door closed. Install a hook-and-eye latch or doorknob cover on the outside of the door. Always close the toilet lid, and consider using a toilet lid latch.
- Never leave a child alone in a bathtub or in the care of another child, even for a moment. Avoid using bath seats or rings. Your baby can slip out of them and be trapped underwater. An adult must be within arm's reach, providing touch supervision at all times.
- Use a rigid, lockable cover on a hot tub, spa, or whirlpool, or fence in all 4 sides as you would for a swimming pool.
- The hottest temperature at the faucet should be no more than 120°F to avoid burns. In many cases, you can adjust your water heater.
- Throw away or tightly cover water or chemical mixtures after use.
- Watch children closely when they are playing near wells, open postholes, or irrigation or drainage ditches. Fill in empty holes or have fences installed to protect your child.
- Learn CPR and know how to get emergency help.

From Your Doctor

The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®





Fun in the Sun:

Keep Your Family Safe

Warm, sunny days are wonderful. It's great to exercise outside, and the sun feels good on your skin. But what feels good can harm you and your family. Read on for information from the American Academy of Pediatrics about how to keep your family safe from the sun's harmful rays.

The Sun and Skin Cancer

The sun gives energy to all living things on earth, but it can also harm us. Its ultraviolet (UV) rays can damage skin and eyes and cause skin cancer. All skin cancers are harmful and some, especially malignant melanoma, can be deadly.

One-quarter of our lifetime sun exposure happens during childhood and adolescence. Since children spend a lot of time outdoors, especially in the summer, it's important to protect them from the sun.

Research shows that 1 or more blistering sunburns as a child or teen can increase the risk of melanoma skin cancer later in life. Sunburns can also be very painful. Too much sun exposure can cause other problems, too, such as

- Dehydration (loss of fluids) and fever
- Damage to skin, such as changes in color and wrinkles
- Cataracts (clouding of eye lens) of the eye
- Damage to the body's immune system

Sun Safety Tips

It's good for children and adults to spend time playing and exercising outdoors, and it's important to do so safely. Follow these simple rules to protect your family from sunburns now and from skin cancer later in life.

- Keep babies younger than 6 months out of direct sunlight. Find shade under a tree, an umbrella, or the stroller canopy.
- When possible, dress yourself and your children in cool, comfortable clothing that covers the body, such as lightweight cotton pants, long-sleeved shirts, and hats.
- Select clothes made with a tight weave; they protect better than clothes with a looser weave. If you're not sure how tight a fabric's weave is, hold it up to see how much light shines through. The less light, the better. Or you can look for protective clothing labeled with an Ultraviolet Protection Factor (UPF).
- Wear a hat with an all-around 3-inch brim to shield the face, ears, and back of the neck.
- Limit your sun exposure between 10:00 am and 4:00 pm when UV rays are strongest.
- Wear sunglasses with at least 99% UV protection. Look for child-sized sunglasses with UV protection for your child.
- Use sunscreen.
- Make sure everyone in your family knows how to protect his or her skin and eyes. Remember to set a good example by practicing sun safety yourself.

Sunscreen

Sunscreen can help protect the skin from sunburn and some skin cancers but only if used correctly. Keep in mind that sunscreen should be used for sun protection, not as a reason to stay in the sun longer.

How to pick sunscreen

- Use a sunscreen that says "broad-spectrum" on the label; that means it will screen out both UVB and UVA rays.
- Use a broad-spectrum sunscreen with a sun protection factor (SPF) of at least 15 (up to SPF 50). An SPF of 15 or 30 should be fine for most people. More research studies are needed to test if sunscreen with more than SPF 50 offers any extra protection.
- If possible, avoid the sunscreen ingredient oxybenzone because of concerns about mild hormonal properties. Remember, though, that it's important to take steps to prevent sunburn, so using any sunscreen is better than not using sunscreen at all.
- For sensitive areas of the body, such as the nose, cheeks, tops of the ears, and shoulders, choose a sunscreen with zinc oxide or titanium dioxide. These products may stay visible on the skin even after you rub them in, and some come in fun colors that children enjoy.

How to apply sunscreen

- Use enough sunscreen to cover all exposed areas, especially the face, nose, ears, feet, hands, and even backs of the knees. Rub it in well.
- Put sunscreen on 15 to 30 minutes before going outdoors. It needs time to absorb into the skin.
- Use sunscreen any time you or your child spends time outdoors. Remember that you can get sunburn even on cloudy days because up to 80% of the sun's UV rays can get through the clouds. Also, UV rays can bounce back from water, sand, snow, and concrete, so make sure you're protected.
- Reapply sunscreen every 2 hours and after swimming, sweating, or drying off with a towel. Because most people use too little sunscreen, make sure to apply a generous amount.

Sunscreen for Babies

For babies younger than 6 months. Use sunscreen on small areas of the body, such as the face, if protective clothing and shade are not available.

For babies older than 6 months. Apply to all areas of the body, but be careful around the eyes. If your baby rubs sunscreen into her eyes, wipe her eyes and hands clean with a damp cloth. If the sunscreen irritates her skin, try a different brand or sunscreen with titanium dioxide or zinc oxide. If a rash develops, talk with your child's doctor.

Sunburns

When to call the doctor

If your baby is younger than 1 year and gets sunburn, call your baby's doctor right away. For older children, call your child's doctor if there is blistering, pain, or fever.

How to soothe sunburn

Here are 5 ways to relieve discomfort from mild sunburn.

1. Give your child water or 100% fruit juice to replace lost fluids.
2. Use cool water to help your child's skin feel better.
3. Give your child pain medicine to relieve painful sunburns. (For a baby 6 months or younger, give acetaminophen. For a child older than 6 months, give either acetaminophen or ibuprofen.)
4. Only use medicated lotions if your child's doctor says it is OK.
5. Keep your child out of the sun until the sunburn is fully healed.

About Indoor Tanning and Sunless Tanning Products

Many teens, especially girls and young women, go to tanning salons because a tan makes them feel more attractive and healthy. But tanning at a salon is dangerous! Like the natural sun, tanning beds give off UV rays that can cause sunburns and skin cancer. Tanning indoors is not safe for anyone—teens or adults!

Sunless tanning lotions, sprays, and airbrush tanning booths are popular too. These products contain a chemical that darkens the skin. The tan usually lasts for several days. However, all sunless tanning products can cause side effects such as skin rashes and irritation. They should also be kept away from the eyes, nose, and mouth. Most of these products do not include sunscreen, so skin is not protected from the real sun. Anyone using a sunless tanner must also use a sunscreen.

Sun Myths

Myth: Only people with light skin can get sunburn.

Fact: People with pale skin or light hair need to be more careful in the sun because they sunburn more easily compared to people with darker skin. However, most people can sunburn no matter what their skin color is. All people need to take steps to protect themselves from the sun's harmful rays.

Myth: A suntan is good for you.

Fact: A "base tan" does not protect you from getting sunburn. In fact, it may increase the chance you'll get sunburn because you may think that you can stay out in the sun longer. A tan is actually a sign of skin damage.

Myth: Only adults can get skin cancer, so putting sunscreen on children is not necessary.

Fact: While most of the people who get skin cancer are older, children, teens, and young adults can get it too. Also, too many sunburns and too much sun exposure over the years can cause not only skin cancer but also skin wrinkles and cataracts of the eye. Skin cancer is the most common form of cancer in the United States. According to the American Cancer Society, there are more than 3.5 million new cases of skin cancer each year.

From Your Doctor



American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



healthychildren.org

Powered by pediatricians. Trusted by parents.
from the American Academy of Pediatrics

The American Academy of Pediatrics (AAP) is an organization of 67,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

The persons whose photographs are depicted in this publication are professional models. They have no relation to the issues discussed. Any characters they are portraying are fictional. The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

© 2019 American Academy of Pediatrics. All rights reserved.

Insect Repellents: What Parents Need to Know

One way to protect your child from biting insects is to use insect repellents. However, it's important that insect repellents are used safely and correctly.

Mosquito, biting fly, and tick bites can make children miserable. While most children have only mild reactions to insect bites, some children can become very sick. Some insects carry dangerous germs such as West Nile virus, Zika virus, Lyme disease bacteria, and Rocky Mountain spotted fever bacteria.

Read on for more information from the American Academy of Pediatrics (AAP) about insect repellents, using repellents safely, and other ways to protect your child from insect bites. Also, read about diseases spread by insects.

About Insect Repellents

Insect repellents prevent bites from biting insects but not stinging insects. Biting insects include mosquitoes, ticks, fleas, chiggers, and biting flies. Stinging insects include bees, hornets, and wasps.

Insect repellents come in many forms, including aerosols, sprays, liquids, creams, and sticks. Some are made from chemicals such as DEET, picaridin, IR3535, and permethrin. Others are made from natural ingredients, such as oil of lemon eucalyptus.

Here are some things to keep in mind.

- Always use an insect repellent registered by the US Environmental Protection Agency (EPA). Visit the US EPA website at www.epa.gov/insect-repellents for more information.
- Always follow the directions.
- Repellents containing 20% to 30% DEET are safe and effective when used as directed for infants and children.
- If you use repellents containing oil of lemon eucalyptus, the Centers for Disease Control and Prevention (CDC) and the AAP recommend that you do not use them on children younger than 3 years. Oil of lemon eucalyptus may cause allergic skin reactions.
- Repellents containing permethrin can be used to treat clothing and gear, but they should not be applied to skin.
- The following types of products offer minimal protection or no protection:
 - Wristbands soaked in chemical repellents
 - Garlic or vitamin B₁ taken by mouth
 - Ultrasonic devices that give off sound waves designed to keep insects away
 - Bird or bat houses
 - Backyard bug zappers, which may attract insects to your yard
 - Essential oils (cedar, cinnamon, citronella, clove, geranium, lemongrass, and others) other than oil of lemon eucalyptus

Tips for Using Insect Repellents Safely

- Read the label and follow all directions and precautions.
- Apply insect repellents only on the outside of your child's clothing and on exposed skin. Never spray an insect repellent directly onto

your child's face. Instead, spray a little on your hands first and then rub it onto your child's face. Avoid your child's eyes and mouth. (Permethrin should not be applied to skin.)

- Do not spray an insect repellent on cuts, wounds, or irritated skin.
- Use just enough of the repellent to cover your child's clothing and exposed skin.
- Avoid reapplying the repellent unless needed. Using more doesn't make the repellent more effective.
- Spray repellents in open areas to avoid breathing them in.
- Apply an insect repellent on young children. Supervise older children when using these products.
- Wash your children's skin with soap and water to remove any repellent when they return indoors, and wash their clothing before they wear it again.
- Do not use products that combine DEET with sunscreen. DEET may make the sun protection factor (SPF) less effective. These products can overexpose your child to DEET because the sunscreen needs to be reapplied often.

Reactions to Insect Repellents

If you suspect that your child is having a reaction, such as a rash, to an insect repellent, stop using the product and wash your child's skin with soap and water. Then call Poison Help at 1-800-222-1222 or your child's doctor. If you go to your child's doctor's office, take the repellent container with you.

Other Ways to Protect Your Child From Insect Bites

Here are ways to help reduce insect bites.

- Tell your child to avoid areas that attract flying insects, such as garbage cans, stagnant pools of water, and flower beds or orchards.
- Dress your child in lightly colored clothing from head to toe, such as long pants, a lightweight long-sleeved shirt, socks, and closed shoes, when you know your child will be exposed to insects. A hat with a wide brim can help keep insects away from your child's face. Mosquito netting may be used over baby carriers or strollers in areas where your baby may be exposed to insects.
- Don't use scented soaps, perfumes, or hair sprays on your child because they may attract insects.
- Keep door and window screens in good repair.
- Check your child's skin at the end of the day if you live in an area where ticks are present and your child has been playing outdoors.

Diseases Spread by Insects and Ticks

Infectious diseases spread by insects are a major cause of illnesses to children and adults worldwide. Here are 4 diseases spread by insects.

West Nile virus is spread by mosquitos.

In the United States, West Nile virus and outbreaks of various types of encephalitis get plenty of media coverage.

Symptoms

- Most cases of West Nile virus infection are mild. People may have no symptoms or mild symptoms such as fever, headache, and body aches.
- Less common symptoms, occurring mostly in older adults, may include a severe headache, a high fever, a stiff neck, confusion, seizures, sensitivity to light, muscle weakness, and loss of consciousness.

Zika virus is primarily spread by mosquitoes.

Zika virus infection during pregnancy can cause fetuses to have a birth defect of the brain called *microcephaly*. Zika virus outbreaks are currently happening in many countries and territories. For up-to-date information about the virus and outbreaks, go to www.cdc.gov/zika.

Symptoms

- Many people won't have symptoms or will have only mild symptoms.
- Symptoms may include fever, rash, joint pain, or red eyes.

Lyme disease is spread by deer ticks.

Deer ticks are tiny, black-brown arachnids about the size of a poppy seed. They are not insects because they have 8 legs (like spiders). Lyme disease is an important health concern in certain regions of the country; the following areas are where most infections occur: Northeast, from Virginia to Maine; north-central states, mostly Wisconsin and Minnesota; and West Coast, particularly northern California.

Symptoms

Often the first and most obvious symptom of Lyme disease is a localized rash that begins as a pink or red circle at the site of tick attachment. This circle expands over time and may become several inches or larger. A classic bull's-eye appearance, with concentric rings, appears in some people. The rash generally occurs 1 to 2 weeks after the tick bite but ranges from 3 days to 30 days.

A rash may occur without any other symptoms or may be associated with

- Headache
- Chills
- Fever
- Fatigue
- Swollen glands, usually in the neck or groin
- Aches and pains in the muscles or joints

If you live in an area of the country endemic to Lyme disease and your child develops a suspicious rash with or without any of these symptoms, call your child's doctor.

Rocky Mountain spotted fever is spread by ticks.

Despite the name, Rocky Mountain spotted fever currently occurs mostly in other regions of the United States, including North and South Carolina, Oklahoma, and Tennessee.

Symptoms

- Most people get a red, dot-like rash that begins on the wrists and ankles and spreads toward the center of the body. The illness may be severe or fatal in some people.
- Other symptoms may include severe headache, fever, muscle aches, nausea, or vomiting.

If your child has been bitten by an insect and shows any of the symptoms of West Nile virus infection, Zika virus infection, Lyme disease, or Rocky Mountain spotted fever, call your child's doctor.

Remember

Children need and love to be outdoors. The chance of your children becoming infected by insects and ticks is quite low. The best way to protect yourself and your children is to follow the guidelines in this publication. If you have any concerns about insect or tick bites, talk with your child's doctor.

From Your Doctor



American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



healthychildren.org

Powered by pediatricians. Trusted by parents.
from the American Academy of Pediatrics

The American Academy of Pediatrics (AAP) is an organization of 67,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of all infants, children, adolescents, and young adults.

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Any websites, brand names, products, or manufacturers are mentioned for informational and identification purposes only and do not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of external resources. Information was current at the time of publication.

© 2019 American Academy of Pediatrics. All rights reserved.



Childproofing Your Home

Children are naturally curious and love to explore. Young children especially like to explore by putting things in their mouths. Before or as soon as children begin crawling or walking, parents and caregivers need to take extra steps to make sure harmful items are out of reach, out of sight, and locked up if possible.

Take a moment to check each room in your home using the Safety Checklist in this publication.

Place a check in the first column next to the item if it is STORED OUT OF REACH OF CHILDREN.

Place a check in the second column next to the item if it is STORED IN A LOCKED CABINET WITH A SAFETY LATCH.

Also keep in mind that children may get into trash containers. Trash containers that contain spoiled food, sharp objects (like discarded razor blades), or batteries should have a child-resistant cover or be kept out of a child's reach. Purses and other bags that hold potential hazards, including medicines, should be kept out of a child's reach too.

Important Reminders

Most poisonings occur when parents or caregivers are home but not paying attention.

- Keep products in original packaging.
- Store in locked cabinets or containers, out of sight and reach of children.
- Install a safety latch that locks when you close the door on child-accessible cabinets.

Detergent in single-use laundry packets is very concentrated and can be toxic. Even a small amount of the detergent can cause serious breathing or stomach problems or eye irritation.

- Never let your children handle or play with the packets. The packets dissolve quickly when in contact with water, wet hands, or saliva. Biting a packet can cause it to burst, shooting detergent into the child's mouth and throat or eyes.
- Remember to seal the container and store it in a locked cabinet after each use. Make sure the container is out of sight and reach of children.
- Adults should follow the instructions on the product label.

Small objects can be choking hazards or harmful if swallowed.

- Check your floors regularly for small objects. This is particularly important if someone in the household has a hobby that involves small items or if there are older children who have small items.
- Make sure battery covers are secure on remote controls, key fobs, musical books, and greeting cards. Store devices that contain small button batteries out of sight and reach of children. Button batteries can cause severe injury or death if ingested.

Safety Checklist

Check your home for...	Stored out of reach of children	Stored in a locked cabinet with a safety latch
Cleaning products including <ul style="list-style-type: none"> • All-purpose cleaners • Bleach • Dishwashing detergent (liquid, powdered, or single-use packets or tablets) • Drain openers and toilet bowl cleaners • Furniture polish • Laundry detergent (liquid, powdered, or single-use packets or tablets) 	☐ ☐ ☐ ☐ ☐ ☐	☐ ☐ ☐ ☐ ☐ ☐
Personal and hygiene products including <ul style="list-style-type: none"> • Nail polish removers • Cosmetics • Mouthwash • Perfume and aftershave 	☐ ☐ ☐ ☐	☐ ☐ ☐ ☐
Items that may be stored in your basement or garage including <ul style="list-style-type: none"> • Antifreeze and windshield wiper fluid • Gasoline, kerosene, and lamp oil • Insecticides and weed killer 	☐ ☐ ☐	☐ ☐ ☐
Medicines <ul style="list-style-type: none"> • Prescription medicines • Over-the-counter medicines • Vitamins and supplements 	☐ ☐ ☐	☐ ☐ ☐
Houseplants Certain houseplants may be harmful. Call Poison Help at 1-800-222-1222 for a list or description of plants to avoid. You may want to do without houseplants for a while or, at the very least, keep all houseplants out of reach.	☐	☐
Small objects including <ul style="list-style-type: none"> • Beads, buttons, coins • Button batteries • Pins • Refrigerator magnets or products and toys with small or loose magnets • Screws 	☐ ☐ ☐ ☐ ☐	☐ ☐ ☐ ☐ ☐
Alcohol Alcohol can be very poisonous to a young child. Remember to empty any unfinished drinks right away.	☐	☐
Nicotine including e-cigarette refills and cigarettes Liquid nicotine e-cigarette refills can be extremely dangerous, even fatal, for a child. Keep all nicotine products, including traditional cigarettes, out of sight and reach of children.	☐	☐

Medicines can be harmful if not taken as directed.

- Purchase and keep medicines in original containers with safety caps.
- Check the label each time you give a child medicine to ensure proper dosage.

From Your Doctor



What to Do in Case of Poisoning

If you find your child with an open or empty container of a dangerous nonfood item, your child may have been poisoned.

1. Stay calm and act quickly.
2. Get the item away from your child. If there is still some in your child's mouth, make your child spit it out or remove it with your fingers. Keep this material along with anything else that might help determine what your child swallowed.
3. Do not make your child vomit because it may cause more damage.

If your child is unconscious, not breathing, or having convulsions or seizures, call 911 or your local emergency number right away.

If your child does not have these symptoms, call Poison Help at 1-800-222-1222. You may be asked for the following information:

- Your name and phone number
- Your child's name, age, and weight
- Any medical conditions your child has
- Any medicine your child is taking
- The name of the item your child swallowed
- The time your child swallowed the item (or when you found your child), and the amount you think was swallowed

If the poison is very dangerous, or if your child is very young, you may be told to take your child to the nearest hospital. If your child is not in danger, the Poison Help staff will tell you what to do to help your child at home.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



healthychildren.org

Powered by pediatricians. Trusted by parents.
from the American Academy of Pediatrics

The American Academy of Pediatrics (AAP) is an organization of 67,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of all infants, children, adolescents, and young adults.

The persons whose photographs are depicted in this publication are professional models. They have no relation to the issues discussed. Any characters they are portraying are fictional. Any websites, brand names, products, or manufacturers are mentioned for informational and identification purposes only and do not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of external resources. Information was current at the time of publication. The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

© 2020 American Academy of Pediatrics. All rights reserved.

Keep Your Family Safe

Fire Safety and Burn Prevention at Home



Fires and burns cause almost 4,000 deaths and about 20,000 hospitalizations every year. Winter is an especially dangerous time, as space heaters, fireplaces, and candles get more use in the home. It is no surprise that fires in the home are more common between December and February. However, you might be surprised at how easy it is to reduce the risk of fire in your home. Follow these suggestions to help keep your home and family safe from fire all year round.

Smoke alarms save lives

Half of home fire deaths are due to fires that happen while people are sleeping. One of the most important steps you can take to protect your family against fire is to install smoke alarms and keep them in good working order. You can buy smoke alarms at most home and hardware stores, and they often cost \$10 or less. Check with your fire department to see if they give out and install free smoke alarms.

- **Install** smoke alarms outside every bedroom or any area where someone sleeps. Also install them in furnace areas. Be sure there is at least 1 alarm on every level of your home, including the basement, or at each end of a mobile home.
- **Place** smoke alarms away from the kitchen and bathroom. False alarms can occur while cooking or even showering.
- **Test** smoke alarms every month by pushing the test button. It is best to use smoke alarms that have long-life batteries, but if you do not, change the batteries at least once a year, such as when you change your clocks in the fall.
- **Replace** smoke alarms every 10 years.
- **Never** paint a smoke alarm.
- **Clean** (dust or vacuum) smoke alarms once a month.
- **Use** smoke alarms equipped with a flashing light and an alarm in homes with children or adults who are hard of hearing or deaf.

Safety around the home

Take a careful look at each room of your home.

Use the following checklists and safety tips to reduce the risk of fire:

- **Do not smoke in your home.** If you do, use deep ashtrays and do not smoke in bed.
- **Make an escape plan.** Practice it every 6 months. Every member of the family should know at least 2 exits from each room and where to meet outside. Make sure doors and windows are easy to open to permit easy escape if needed.
- **Check electrical cords.** Replace any electrical cords that are worn, frayed, or damaged. Never overload outlets. Avoid running electrical cords under carpets or furniture because they can overheat and start a fire.

- **Consider installing an automatic home fire sprinkler system.** They are now practical for many homes.
- **Ask your local fire department to make sure woodstoves in your home are safely vented.** They usually cannot be installed safely in mobile homes.
- **Avoid using kerosene heaters and electric space heaters.** If electric space heaters must be used, keep them away from clothing, bedding, and curtains, and unplug them at night. Kerosene heaters give off carbon monoxide and should not be used in enclosed spaces, such as inside your home.

Bedrooms

- ☐ Check the labels of your child's pajamas. Children should always wear flame-retardant and/or close-fitting sleepwear.
- ☐ If a bedroom is on an upper floor, make sure there is a safe way to reach the ground, such as an escape ladder that will not burn.
 - ♦ *Never smoke in bed or when you are drowsy or have been drinking. Tobacco and smoking products, matches, and lighters are the most common cause of fatal fires in the home.*

Living and family rooms

Make sure all matches, lighters, and ashtrays are out of your child's sight and reach. Better yet, keep them in a locked cabinet.

- ☐ Use large, deep ashtrays that won't tip over, and empty them often. Fill ashtrays with water before dumping ashes in the wastebasket.
- ☐ Give space heaters plenty of space. Keep heaters at least 3 feet from anything that might burn, like clothes, curtains, and furniture. Always turn space heaters off and unplug them when you go to bed or leave the home.
- ☐ Have fireplaces and chimneys cleaned and inspected once a year.
- ☐ Use a metal screen or glass doors in front of the fireplace.
 - ♦ *Never leave a room unattended when candles, heaters, or fireplaces are in use.*

Kitchen

- ☐ Keep your stove and oven clean and free of anything that could catch fire. Do not place pot holders, curtains, or towels near the burners.
- ☐ Install a portable fire extinguisher in the kitchen, high on a wall, and near an exit. (Choose a multipurpose, dry chemical extinguisher.) Adults should know how to use it properly when the fire is small and contained, such as in a trash can. Call your fire department for information on how to use fire extinguishers.
 - ♦ *Never leave cooking food unattended.*

- ◆ *Never pour water on a grease fire.*
- ◆ *If a fire starts in your oven, keep the oven door closed, turn off the oven, and call the fire department.*

Garage, storage area, and basement

- Have your furnace inspected at least once a year.
- Do not store anything near a heater or furnace. Keep the area free of clutter.
- Clean your dryer's lint filter after every use. Lint buildup can start a fire.
- Check to make sure paint and other flammable liquids are stored in their original containers, with tight-fitting lids. Store them in a locked cabinet if possible, out of your child's reach, and away from appliances, heaters, pilot lights, and other sources of heat or flame.
- Never use flammable liquids near a gas water heater.
- Store gasoline, propane, and kerosene outside the home in a shed or detached garage. Keep them tightly sealed and labeled in approved safety containers.
 - ◆ *Gasoline should be used only as a motor fuel, never as a cleaning agent.*
 - ◆ *Never smoke near flammable liquids.*

Outdoors

- Move barbecue grills away from trees, bushes, shrubs, or anything that could catch fire. Never use grills indoors, on a porch, or on a balcony.
- Place a barrier around open fires, fire pits, or campfires. Never leave a child alone around the fire. Always be sure to put the fire out completely before leaving or going to sleep.
- Create a "fire break" around your home. Make sure woodpiles, dead leaves, pine needles, and debris are removed or kept as far away from the home as possible.
 - ◆ *Do not start lawnmowers, snow blowers, or motorcycles near gasoline fumes. Let motors cool off before adding fuel.*
 - ◆ *Be very careful with barbecue grills. Never use gasoline to start the fire. Do not add charcoal lighter fluid once the fire has started.*

Know what to do in a fire

- **Test any closed doors with the back of your hand for heat.** Do not open the door if you feel heat or see smoke. Close all doors as you leave each room to keep the fire from spreading.
- **Crawl low under smoke.** Choose the safest exit. If you must escape through a smoky area, remember that cleaner air is always near the floor. Teach your child to crawl on her hands and knees, keeping her head less than 2 feet above the floor, as she makes her way to the nearest exit.
- **Don't stop. Don't go back.** In case of fire, do not try to rescue pets or possessions. Once you are out, do not go back in for any reason. Firefighters have the best chance of rescuing people who are trapped. Let firefighters know right away if anyone is missing.
- **If you get trapped by smoke or flames,** close all doors. Stuff towels or clothing under the doors to keep out smoke. Cover your nose and mouth with a damp cloth to protect your lungs. If there is no phone in the room, wait at a window and signal for help with a light-colored cloth or flashlight.
- **Stop, drop, and roll! Cool and call.** Make sure your child knows what to do if her clothes catch fire.
 - Stop!*—Do not run.

Drop!—Drop to the ground right where you are.

Roll!—Roll over and over to put out the flames. Cover your face with your hands.

Cool!—Cool the burned area with water.

Call!—Call for help.

Fire and children

A child's curiosity about fire is natural and in most cases is no cause for concern. However, when a child begins to use fire as a weapon, it can be very dangerous. If you suspect that your child is setting even very small fires, address the problem right away. Talk with your pediatrician, who can suggest ways to help.

Use the following tips when talking with your child about preventing fires:

- Teach your child that matches and lighters are tools for grown-ups only.
- Older children should be taught to use fire properly, and **only** with an adult present.

For your sitters

When you are away from home and someone else cares for your children, make sure that your children and the sitter will be just as safe as when you are there.

- Let your sitter know about your family's escape plan.
- Remind sitters never to leave the children alone.
- Remind sitters that you do not allow smoking in or around your home.

Leave emergency information near the phone. Include the local fire department phone number, your full home address and phone number, and a neighbor's name and phone number.

Burn prevention

Most burn injuries happen in the home. For a young child, many places in the home can be dangerous.

Hot bathwater, radiators, and even food that is too hot can cause burns. The following are tips to help prevent your child from getting burned:

- Keep matches, lighters, and ashtrays out of the reach of children.
- Cover all unused electrical outlets with plastic plugs or other types of outlet covers.
- Do not allow your child to play close to fireplaces, radiators, or space heaters.
- Replace all frayed, broken, or worn electrical cords.
- Never leave barbecue grills unattended.
- Teach your children that irons, curling irons, grills, radiators, and ovens can get very hot and are dangerous to touch or play near. Never leave these items unattended. Unplug and put away all appliances after using them.
- Keep electrical cords from hanging down where children can pull on them or chew on them. Mouth burns can result from chewing on a live extension cord or on a poorly insulated wire.

Kitchen concerns

- Never leave a child alone in the kitchen when food is cooking.
- Enforce a "kid-free" zone at least 3 feet around the oven or stove while you are cooking. Use a playpen, high chair, or other stationary device to keep your child from getting too close.

- Never leave a hot oven door open.
- Use back burners if possible. When using front burners, turn pot handles rearward. Never let them stick out where a child could grab them.
- Do not leave spoons or other utensils in pots while cooking.
- Turn off burners and ovens when they are not being used.
- Do not use wet pot holders because they may cause steam burns.
- Carefully place (not toss) wet foods into a deep fryer or frying pan containing grease. The reaction between hot oil and water causes splatter.
- Remove pot lids carefully to avoid being burned by steam. Remember, steam is hotter than boiling water.
- In case of a small pan fire, carefully slide a lid over the pan to smother the flames, turn off the burner, and wait for the pan to cool completely.
- Never carry your child and hot liquids at the same time.
- Never leave hot liquid, like a cup of coffee, where children can reach it. Don't forget that a child can get burned from hot liquids by pulling on hanging tablecloths.
- Wear tight-fitting or rolled-up sleeves when cooking to reduce the risk of your clothes catching on fire.
- In microwave ovens, use only containers that are made for microwaves. Test microwaved food for heat and steam before giving it to your child. (Never warm a bottle of milk or formula in the microwave oven. It can heat the liquid unevenly and burn your child.)
- Avoid letting appliance cords hang over the sides of countertops, where children could pull on them.
- Do not use mobile baby walkers. They allow your child to move quickly before he knows how to use this mobility safely. It may allow him to gain access to hot liquids, appliance cords, and hot surfaces.

Hot water

- The hottest water temperature at the faucet should be no higher than 120°F to prevent scalding. In many cases, you can adjust your water heater to prevent exceeding this temperature.
- When using tap water, always turn on the cold water first, then add hot. When finished, turn off the hot water first.
- Test the temperature of bathwater with your forearm or the back of your hand before placing your child in the water.
- Use a cool-mist vaporizer instead of a hot-water vaporizer. Hot-water vaporizers can cause steam burns or can spill on your child.
- Never leave children alone in the bathroom for any reason. They are at risk of burns and drowning.

First aid for burns

For severe burns, call 911 or your local emergency number right away. Until help arrives, follow these steps.

1. Cool the burn.

For 1st and 2nd degree burns, cool the burned area with cool running water for a few minutes. This helps stop the burning process, numbs the pain, and prevents or reduces swelling. Do not use ice on a burn. It may delay healing. Also, do not rub a burn; it can increase blistering.

For 3rd degree burns, cool the burn with wet, sterile dressings until help arrives.

2. Remove burned clothing.

Lay the person flat on her back and take off the burned clothing that isn't stuck to the skin. Remove any jewelry or tight-fitting clothing from around the burned area before swelling begins. If possible, elevate the injured area.

3. Cover the burn.

After the burn has cooled, cover it loosely with a dry bandage or clean cloth. Do not break any blisters. This could allow bacteria into the wound. Never put grease (including butter or medical ointments) on the burn. Grease holds in heat, which may make the burn worse. It also makes the burn harder to examine by medical personnel later.

4. Keep the child from losing body heat.

Keep the person's body temperature normal. Cover unburned areas with a dry blanket.

Fire drills—be prepared!

Even young children (3 and older) can begin to learn what to do in case of a fire.

Install at least 1 smoke alarm on every level of your home.

Have an escape plan and practice it with your family. This will help you and your family reach safety when it counts. When a fire occurs, there will be no time for planning an escape.

Draw a floor plan of your home. Discuss with your family 2 ways to exit every room. Make sure everyone knows how to get out and that doors and windows can be easily opened to permit escape.

If you live in an apartment building, never use an elevator during a fire. Use the stairs!

Agree on a meeting place. Choose a spot outside your home near a tree, street corner or fence where everyone can meet after escaping. Teach your children that the sound of a smoke alarm means to go outside right away to the chosen place.

Know how to call the fire department. The fire department should be called from outside using a portable phone or from a neighbor's home. Whether the number is 911 or a regular phone number, everyone in the family should know it by heart. Make sure your children know your home address too. Teach your children that firefighters are friends and never to hide from them.

Practice, practice, practice. Practice your exit drill at least twice a year. Remember that fire drills are not a race. Get out quickly, but calmly and carefully. Try practicing realistic situations. Pretend that some exits or doorways are blocked or that the lights are out. The more prepared your family is, the better your chances of surviving a fire.

Note: Parents of very young children or children with special needs should have a safety plan that fits their child's needs and abilities. For example, a child who is hard of hearing or deaf may need a smoke alarm with a flashing strobe-light feature. Parents with children younger than 5 years must plan on an adult rescuing them in the case of a house fire; they are too young to be able to reliably rescue themselves.

Different degrees of burns

Following are the 4 different levels of burns and the symptoms of each:

1st degree burns are minor and heal quickly. Symptoms are redness, tenderness, and soreness (like most sunburns).

2nd degree burns are serious injuries. First aid and medical treatment should be given as soon as possible. Symptoms are blistering (like a severe sunburn), pain, and swelling.

3rd degree burns (also called full-thickness burns) are severe injuries. Medical treatment is needed right away. Symptoms are white, brown, or charred tissue often surrounded by blistered areas. There may be little or no pain at first.

4th degree burns are severe injuries that involve skin, muscle, and bone. These often occur with electrical burns and may be more severe than they appear. They may cause serious complications and should be treated by a doctor right away.

Call your pediatrician if your child suffers anything more than a minor burn. ALL electrical burns and any burn on the hand, foot, face, genitals, or over a joint worse than 1st degree should receive medical attention right away.

From your doctor

Adapted from material provided by the National Fire Protection Association (NFPA). For more information, call 617/770-3000, or visit the NFPA Web site at www.nfpa.org or its family Web site at www.sparky.org.

Please note: Listing of resources does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this publication. Phone numbers and Web site addresses are as current as possible, but may change at any time.

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The American Academy of Pediatrics is an organization of 60,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

American Academy of Pediatrics
Web site—www.aap.org

Copyright © 2007
American Academy of Pediatrics, Updated 4/07

Your child needs vaccines as they grow!

2025 Recommended Immunizations for Birth Through 6 Years Old

Want to learn more?
Scan this QR code to find out which
vaccines your child might need. Or visit
www2.cdc.gov/vaccines/childquiz/



VACCINE OR PREVENTIVE ANTIBODY	BIRTH	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS	7 MONTHS	8 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	19 MONTHS	20–23 MONTHS	2–3 YEARS	4–6 YEARS
RSV antibody	Depends on mother's RSV vaccine status						Depends on child's health status							
Hepatitis B	Dose 1	Dose 2			Dose 3									
Rotavirus			Dose 1	Dose 2	Dose 3									
DTaP			Dose 1	Dose 2	Dose 3				Dose 4					Dose 5
Hib			Dose 1	Dose 2	Dose 3			Dose 4						
Pneumococcal			Dose 1	Dose 2	Dose 3			Dose 4						
Polio			Dose 1	Dose 2	Dose 3								Dose 4	
COVID-19					At least 1 dose of the current COVID-19 vaccine									
Influenza/Flu					Every year. Two doses for some children									
MMR								Dose 1						Dose 2
Chickenpox								Dose 1						Dose 2
Hepatitis A								2 doses separated by 6 months						

KEY

- ALL children should be immunized at this age
- SOME children should get this dose of vaccine or preventive antibody at this age

Talk to your child's health care provider for more guidance if:

1. Your child has any medical condition that puts them at higher risk for infection.
2. Your child is traveling outside the United States. Visit wwwnc.cdc.gov/travel for more information.
3. Your child misses a vaccine recommended for their age.

What diseases do these vaccines protect against?

BIRTH–6 YEARS OLD

VACCINE-PREVENTABLE DISEASE	DISEASE COMPLICATIONS
RSV (Respiratory syncytial virus) Contagious viral infection of the nose, throat, and sometimes lungs; spread through air and direct contact	Infection of the lungs (pneumonia) and small airways of the lungs; especially dangerous for infants and young children
Hepatitis B Contagious viral infection of the liver; spread through contact with infected body fluids such as blood or semen	Chronic liver infection, liver failure, liver cancer, death
Rotavirus Contagious viral infection of the gut; spread through the mouth from hands and food contaminated with stool	Severe diarrhea, dehydration, death
Diphtheria* Illness caused by a toxin produced by bacteria that infects the nose, throat, and sometimes skin	Swelling of the heart muscle, heart failure, coma, paralysis, death
Pertussis (Whooping Cough)* Contagious bacterial infection of the lungs and airway; spread through air and direct contact	Infection of the lungs (pneumonia), death; especially dangerous for babies
Tetanus (Lockjaw)* Bacterial infection of brain and nerves caused by spores found in soil and dust everywhere; spores enter the body through wounds or broken skin	Seizures, broken bones, difficulty breathing, death
Hib (Haemophilus influenzae type b) Contagious bacterial infection of the lungs, brain and spinal cord, or bloodstream; spread through air and direct contact	Depends on the part of the body infected, but can include brain damage, hearing loss, loss of arm or leg, death
Pneumococcal Bacterial infections of ears, sinuses, lungs, or bloodstream; spread through direct contact with respiratory droplets like saliva or mucus	Depends on the part of the body infected, but can include infection of the lungs (pneumonia), blood poisoning, infection of the lining of the brain and spinal cord, death
Polio Contagious viral infection of nerves and brain; spread through the mouth from stool on contaminated hands, food or liquid, and by air and direct contact	Paralysis, death
COVID-19 Contagious viral infection of the nose, throat, or lungs; may feel like a cold or flu. Spread through air and direct contact	Infection of the lungs (pneumonia); blood clots; liver, heart or kidney damage; long COVID; death
Influenza (Flu) Contagious viral infection of the nose, throat, and sometimes lungs; spread through air and direct contact	Infection of the lungs (pneumonia), sinus and ear infections, worsening of underlying heart or lung conditions, death
Measles (Rubeola)† Contagious viral infection that causes high fever, cough, red eyes, runny nose, and rash; spread through air and direct contact	Brain swelling, infection of the lungs (pneumonia), death
Mumps† Contagious viral infection that causes fever, tiredness, swollen cheeks, and tender swollen jaw; spread through air and direct contact	Brain swelling, painful and swollen testicles or ovaries, deafness, death
Rubella (German Measles)† Contagious viral infection that causes low-grade fever, sore throat, and rash; spread through air and direct contact	Very dangerous in pregnant women; can cause miscarriage or stillbirth, premature delivery, severe birth defects
Chickenpox (Varicella) Contagious viral infection that causes fever, headache, and an itchy, blistering rash; spread through air and direct contact	Infected sores, brain swelling, infection of the lungs (pneumonia), death
Hepatitis A Contagious viral infection of the liver; spread by contaminated food or drink or close contact with an infected person	Liver failure, death

*DTaP protects against tetanus, diphtheria, and pertussis

†MMR protects against measles, mumps, and rubella

DTaP (Diphtheria, Tetanus, Pertussis) Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

DTaP vaccine can prevent **diphtheria, tetanus, and pertussis**.

Diphtheria and pertussis spread from person to person. Tetanus enters the body through cuts or wounds.

- **DIPHTHERIA (D)** can lead to difficulty breathing, heart failure, paralysis, or death.
- **TETANUS (T)** causes painful stiffening of the muscles. Tetanus can lead to serious health problems, including being unable to open the mouth, having trouble swallowing and breathing, or death.
- **PERTUSSIS (aP)**, also known as “whooping cough,” can cause uncontrollable, violent coughing that makes it hard to breathe, eat, or drink. Pertussis can be extremely serious especially in babies and young children, causing pneumonia, convulsions, brain damage, or death. In teens and adults, it can cause weight loss, loss of bladder control, passing out, and rib fractures from severe coughing.

2. DTaP vaccine

DTaP is only for children younger than 7 years old. Different vaccines against tetanus, diphtheria, and pertussis (Tdap and Td) are available for older children, adolescents, and adults.

It is recommended that children receive 5 doses of DTaP, usually at the following ages:

- 2 months
- 4 months
- 6 months
- 15–18 months
- 4–6 years

DTaP may be given as a stand-alone vaccine, or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

DTaP may be given at the same time as other vaccines.

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any vaccine that protects against tetanus, diphtheria, or pertussis**, or has any **severe, life-threatening allergies**
- Has had a **coma, decreased level of consciousness, or prolonged seizures within 7 days after a previous dose of any pertussis vaccine (DTP or DTaP)**
- Has **seizures or another nervous system problem**
- Has ever had **Guillain-Barré Syndrome** (also called “GBS”)
- Has had **severe pain or swelling after a previous dose of any vaccine that protects against tetanus or diphtheria**

In some cases, your child’s health care provider may decide to postpone DTaP vaccination until a future visit.

Children with minor illnesses, such as a cold, may be vaccinated. Children who are moderately or severely ill should usually wait until they recover before getting DTaP vaccine.

Your child’s health care provider can give you more information.



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

4. Risks of a vaccine reaction

- Soreness or swelling where the shot was given, fever, fussiness, feeling tired, loss of appetite, and vomiting sometimes happen after DTaP vaccination.
- More serious reactions, such as seizures, non-stop crying for 3 hours or more, or high fever (over 105°F) after DTaP vaccination happen much less often. Rarely, vaccination is followed by swelling of the entire arm or leg, especially in older children when they receive their fourth or fifth dose.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636** (**1-800-CDC-INFO**) or
 - Visit CDC's website at www.cdc.gov/vaccines.



Pneumococcal Conjugate Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Pneumococcal conjugate vaccine can prevent pneumococcal disease.

Pneumococcal disease refers to any illness caused by pneumococcal bacteria. These bacteria can cause many types of illnesses, including pneumonia, which is an infection of the lungs. Pneumococcal bacteria are one of the most common causes of pneumonia.

Besides pneumonia, pneumococcal bacteria can also cause:

- Ear infections
- Sinus infections
- Meningitis (infection of the tissue covering the brain and spinal cord)
- Bacteremia (infection of the blood)

Anyone can get pneumococcal disease, but children under 2 years old, people with certain medical conditions or other risk factors, and adults 65 years or older are at the highest risk.

Most pneumococcal infections are mild. However, some can result in long-term problems, such as brain damage or hearing loss. Meningitis, bacteremia, and pneumonia caused by pneumococcal disease can be fatal.

2. Pneumococcal conjugate vaccine

Pneumococcal conjugate vaccine helps protect against bacteria that cause pneumococcal disease. There are three pneumococcal conjugate vaccines (PCV13, PCV15, and PCV20). The different vaccines are recommended for different people based on age and medical status. Your health care provider can help you determine which type of pneumococcal conjugate vaccine, and how many doses, you should receive.

Infants and young children usually need 4 doses of pneumococcal conjugate vaccine. These doses are recommended at 2, 4, 6, and 12–15 months of age.

Older children and adolescents might need pneumococcal conjugate vaccine depending on their age and medical conditions or other risk factors if they did not receive the recommended doses as infants or young children.

Adults 19 through 64 years old with certain medical conditions or other risk factors who have not already received pneumococcal conjugate vaccine should receive pneumococcal conjugate vaccine.

Adults 65 years or older who have not previously received pneumococcal conjugate vaccine should receive pneumococcal conjugate vaccine.

Some people with certain medical conditions are also recommended to receive pneumococcal polysaccharide vaccine (a different type of pneumococcal vaccine, known as PPSV23). Some adults who have previously received a pneumococcal conjugate vaccine may be recommended to receive another pneumococcal conjugate vaccine.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any type of pneumococcal conjugate vaccine (PCV13, PCV15, PCV20, or an earlier pneumococcal conjugate vaccine known as PCV7), or to any vaccine containing diphtheria toxoid** (for example, DTaP), or has any **severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone pneumococcal conjugate vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover.

Your health care provider can give you more information.

4. Risks of a vaccine reaction

- Redness, swelling, pain, or tenderness where the shot is given, and fever, loss of appetite, fussiness (irritability), feeling tired, headache, muscle aches, joint pain, and chills can happen after pneumococcal conjugate vaccination.

Young children may be at increased risk for seizures caused by fever after a pneumococcal conjugate vaccine if it is administered at the same time as inactivated influenza vaccine. Ask your health care provider for more information.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636** (**1-800-CDC-INFO**) or
 - Visit CDC's website at www.cdc.gov/vaccines.



Rotavirus Vaccine:

What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Rotavirus vaccine can prevent **rotavirus disease**.

Rotavirus commonly causes severe, watery diarrhea, mostly in babies and young children. Vomiting and fever are also common in babies with rotavirus. Children may become dehydrated and need to be hospitalized and can even die.

2. Rotavirus vaccine

Rotavirus vaccine is administered by putting drops in the child's mouth. Babies should get 2 or 3 doses of rotavirus vaccine, depending on the brand of vaccine used.

- The first dose must be administered before 15 weeks of age.
- The last dose must be administered by 8 months of age.

Almost all babies who get rotavirus vaccine will be protected from severe rotavirus diarrhea.

Another virus called "porcine circovirus" can be found in one brand of rotavirus vaccine (Rotarix). This virus does not infect people, and there is no known safety risk.

Rotavirus vaccine may be given at the same time as other vaccines.

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of rotavirus vaccine**, or has any **severe, life-threatening allergies**
- Has a **weakened immune system**
- Has **severe combined immunodeficiency (SCID)**
- Has had a type of bowel blockage called "**intussusception**"

In some cases, your child's health care provider may decide to postpone rotavirus vaccination until a future visit.

Infants with minor illnesses, such as a cold, may be vaccinated. Infants who are moderately or severely ill should usually wait until they recover before getting rotavirus vaccine.

Your child's health care provider can give you more information.

4. Risks of a vaccine reaction

- Irritability or mild, temporary diarrhea or vomiting can happen after rotavirus vaccine.

Intussusception is a type of bowel blockage that is treated in a hospital and could require surgery. It happens naturally in some infants every year in the United States, and usually there is no known reason for it. There is also a small risk of intussusception from rotavirus vaccination, usually within a week after the first or second vaccine dose. This additional risk is estimated to range from about 1 in 20,000 U.S. infants to 1 in 100,000 U.S. infants who get rotavirus vaccine. Your health care provider can give you more information.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

5. What if there is a serious problem?

For intussusception, look for signs of stomach pain along with severe crying. Early on, these episodes could last just a few minutes and come and go several times in an hour. Babies might pull their legs up to their chest. Your baby might also vomit several times or have blood in the stool, or could appear weak or very irritable. These signs would usually happen during the first week after the first or second dose of rotavirus vaccine, but look for them any time after vaccination. If you think your baby has intussusception, contact a health care provider right away. If you can't reach your health care provider, take your baby to a hospital. Tell them when your baby got rotavirus vaccine.

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636** (**1-800-CDC-INFO**) or
 - Visit CDC's website at www.cdc.gov/vaccines.



IMMUNIZATION INFORMATION STATEMENT

Respiratory Syncytial Virus (RSV) Preventive Antibody: *What You Need to Know*

Why get immunized with a RSV preventive antibody?

A respiratory syncytial virus (RSV) preventive antibody can prevent severe lung disease caused by RSV.

RSV is a common respiratory virus that usually causes mild, cold-like symptoms but can also affect the lungs. Symptoms of RSV infection may include runny nose, decrease in appetite, coughing, sneezing, fever, or wheezing.

Anyone can become infected by RSV, and almost all children get an RSV infection by the time they are 2 years old. While most children recover from an RSV infection in a week or two, RSV infection can be dangerous for infants and some young children, causing difficulty breathing, low oxygen levels, and dehydration. In the United States, RSV is the most common cause of bronchiolitis (inflammation of the small airways in the lungs) and pneumonia (infection of the lungs) in children younger than 1 year of age. Children who get sick from RSV may need to be hospitalized, and some might even die.

RSV Preventive Antibodies

The RSV preventive antibody (generic name nirsevimab, trade name Beyfortus) is a shot that prevents severe RSV disease in infants and young children. Antibodies are proteins that the body's immune system uses to fight off harmful germs. Like traditional vaccines, preventive antibodies are immunizations that provide protection against a specific pathogen. While both are immunizations, the way they provide immunity is different. Nirsevimab is an immunization that provides antibodies directly to the recipient. Traditional vaccines are immunizations that stimulate the recipient's immune system to produce antibodies.

Infants born during the RSV season (typically fall through spring) should receive a single dose of the RSV Immunization within 1 week after birth. Most infants whose mothers got the RSV vaccine don't need to get nirsevimab, too. Both protect infants from severe RSV by providing antibodies, either from the mother to the infant or directly to the infant. Most infants will likely only need protection from either the maternal RSV vaccine or nirsevimab (not both). However, there may be some situations in which nirsevimab would be recommended for an infant after the mother received an RSV vaccine.

Infants born outside of the RSV season who are younger than 8 months should receive a single dose of the RSV Immunization shortly before their first RSV season (typically the fall), but infants who are younger than 8 months who have not yet received a dose may receive a dose at any time during the season.

Some infants and young children who are at increased risk for severe RSV disease may need a single dose of the RSV antibody before or during their second RSV season.

RSV preventive antibodies can be given at the same time as vaccines routinely recommended for infants and young children.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

Talk with your health care provider

Tell your health care provider if the person getting the preventive antibody has a:

- History of serious allergic reactions to an RSV preventive antibody (nirsevimab) or any of its components,
- Bleeding disorder, or
- Moderate or severe acute illness.

In some cases, your child's health care provider may decide to postpone giving RSV preventive antibodies until a future visit.

People who have a minor illness, such as a cold, can safely receive an RSV preventive antibody. People who are moderately or severely ill should usually wait until they recover.

Your health care provider can give you more information.

Risks of a reaction to RSV preventive antibodies

After getting an RSV preventive antibody, your child might have temporary pain, redness, swelling where the injection was given, or a rash.

As with any medicine, there is a very remote chance that RSV Immunization could cause a severe allergic reaction, other serious injury, or death.

An allergic reaction could occur after your child leaves the hospital or clinic. If you see signs of a severe allergic reaction (for example, hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call 9-1-1 and get your child to the nearest hospital.

Call your health care provider if you see any other symptoms that concern you.

What if there is a serious problem?

If your child got an RSV preventive antibody without getting a vaccine at the same time, and you suspect an adverse reaction, you or your health care provider can submit a report through <https://www.fda.gov/medwatch> or by phone at 1-800-FDA-1088.

If your child got an RSV preventive antibody and a vaccine at the same time and you suspect an adverse reaction, you or your health care provider should report it to the [Vaccine Adverse Event Reporting System \(VAERS\)](https://vaers.hhs.gov/) <https://vaers.hhs.gov/> or call [1-800-822-7967](https://vaers.hhs.gov/). In your report, note that your child got an RSV Immunization along with a vaccine.

Note: MedWatch and VAERS are only for reporting reactions. MedWatch and VAERS staff members do not give medical advice.

How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit U.S. Food and Drug Administration website at [Drugs@FDA: FDA-Approved Drugs](https://www.fda.gov/drugs).
- Contact the Centers for Disease Control and Prevention (CDC):
 - o Call 1-800-232-4636 (1-800-CDC-INFO) or
 - o [Visit the CDC website https://www.cdc.gov/rsv/about/prevention.html](https://www.cdc.gov/rsv/about/prevention.html)

