

DODINE GROUP U12 FUNGICIDE

Sequence

FUNGICIDE

For control of certain diseases of Almonds, Apples, Apricots[*], Bananas[*], Beech nuts[*], Brazil nuts[*], Butternuts[*], Cashews[*], Cherries, Chestnuts[*], Chinquapins[*], Filberts[*], Hickory nuts[*], Macadamia nuts[*], Nectarines, Peaches, Peanuts[*], Pears, Pecans[*], Plums[*], Plums-chickasaw[*], Plums-damson[*], Plums-Japanese[*], Plumcots[*], Prune plums[*], and Walnuts.
[*Not registered for use by California.]

ACTIVE INGREDIENT:	WT. BY %
Dodine (n-dodecylguanidine acetate)	39.6%**
OTHER INGREDIENTS:	60.4%
TOTAL:	100.0%

**Equivalent to 3.4 pounds Dodine per gallon

KEEP OUT OF REACH OF CHILDREN DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

FIRST AID - IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to by a poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBERS - For general information about this product, contact the National Pesticides Information Center (NPIC) at **1-800-858-7378**, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>. Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at **1-800-222-1222**.

NOTE TO PHYSICIAN - Probable mucosal damage may contraindicate the usage of gastric lavage.

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 82633-104-83529

EPA Est. No. **AG** 72159-GA-001; **MC** 89332-GA-001; **SC** 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER/PELIGRO

Corrosive. Causes irreversible eye damage. Causes skin irritation. **DO NOT** get in eyes, on skin, or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid breathing spray mist or vapor. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate or viton.

Applicators mixers, loaders, applicators, and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material (barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, and/or viton \geq 14 mils)
- Goggles or face shield
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron for mixers and loaders

Mixers, loaders, and applicators for mechanically pressurized handguns to bananas, olives, Pome Fruits – Crop Group 11-10, Stone Fruits – Crop Group 12-12, and Tree Nuts – Crop Group 14-12 must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Mixers and loaders for aerial applications to bananas, olives, Pome Fruits – Crop Group 11-10, Stone Fruits – Crop Group 12-12, and Tree Nuts – Crop Group 14-12 must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

See **ENGINEERING CONTROLS** for additional requirements.

USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.249(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Surface Water and Erosion Control Statement: This product may contaminate water through runoff and has the potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated including plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants,
- Chemical-resistant gloves made of any waterproof material, chemical-resistant footwear plus socks,
- Protective eyewear, and
- Chemical-resistant headgear (if overhead exposure).

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

USE RESTRICTIONS

- **DO NOT** use this product for any crop or any pest not specifically listed on this label. For crops and pests listed on the label, follow closely the listed application timings.
- **DO NOT** apply at any application timing not listed on the label.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply in greenhouses.
- **DO NOT** graze livestock in treated fields, orchards or treated groves.

FUNGICIDE RESISTANCE MANAGEMENT

For resistance management, **Sequence** contains a Group U12 fungicide. Any fungal population may contain individuals naturally resistant to **Sequence** and other Group U12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides/bactericides are used repeatedly in the same fields. Appropriate resistance management strategies must be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of **Sequence** or other Group U12 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Sharda USA LLC at <http://www.shardausa.com/>. You can also contact your pesticide distributor or university extension specialist to report resistance.

DILUTE AND CONCENTRATE SPRAYS

Sequence may be used effectively in dilute and concentrate sprays when applied properly to deliver the labeled rate (pints per acre). Trees must be covered uniformly. **DO NOT** excessively drench fruit and foliage with dilute sprays. Avoid heavy deposits when using concentrate sprays.

Use **Sequence** in a minimum of 15 gallons of water per acre for all ground applications, and in a minimum of 10 gallons of water per acre for all air applications.

INSTRUCTIONS FOR MIXING

Add **Sequence** directly to the spray tank when half filled with water. Keep agitator running during filling and spraying.

COMPATIBILITY

Experience indicates that **Sequence** can be used effectively at specified dosages with most of the commonly used fungicides, herbicides, insecticides and acaricides. However, some incompatibilities were identified, especially with some copper, oil, sulfur, foliar nutrients and chlorpyrifos formulations. Therefore, we always advise testing the compatibility of the products used in combination with **Sequence** at small scale before spraying or to ask your local distributor or consultant about potential incompatibility.

Observe all precautions, restrictions and limitations on labeling of all products used in mixtures. Follow the most restrictive label limitations and precautions. No label dosage rates may be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity), and method of application (e.g., ground, aerial airstream) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product.

WIND SPEED

DO NOT apply at wind speeds greater than 10 mph at the application site.

DROPLET SIZE

Apply as a medium or coarser spray (ASAE standard 572), except when making orchard airblast applications. For orchard airblast applications, apply at the largest droplet size that provides sufficient coverage and control.

GROUND APPLICATION

Release Height - **DO NOT** release spray at a height greater than 4 feet above the ground or crop canopy.

AIRBLAST

Sprays must be directed into the crop canopy. Turn off outward pointing nozzles at row ends and when spraying outer rows.

AERIAL APPLICATION

- **Release Height - DO NOT** release spray at a height greater than 10 feet above the ground or crop canopy.
- **Boom Length** - The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- **Swath Adjustment** - When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for the displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Leave at least one swath unsprayed at the downwind edge of the treated field.

CROP-SPECIFIC INSTRUCTIONS

**ALMONDS, BEECH NUTS[*], CHESTNUTS[*], AND CHINQUAPINS[*]
[*Not registered for use by California]**

DISEASE	APPLICATION DIRECTIONS	USE RATE
Brown rot blossom blight – (<i>Monilinia laxa</i> / <i>Monilinia fructicola</i>)	For disease suppression at pink bud, full bloom, and petal fall.	3 pints/A (1.3 lbs. a.i./acre)
Scab (<i>Venturia carpophila</i> anamorph, <i>Caldosporium carpophilum</i>)	Use at petal fall. Repeat application 2 and 5 weeks later.	
Shot hole (<i>Wilsonomyces carpophilus</i>)	Use at leaf emergence (bloom through petal fall). Repeat application 2 and 5 weeks later.	
Anthracnose (<i>Gnomonia leptostyla</i>)	Use when first leaves unfold. Repeat application at 10- to 14-day intervals.	

RESTRICTIONS:

- **DO NOT** apply more than 9 pints of **Sequence** per acre per year (3.9 lbs. a.i./acre/year).
- **DO NOT** apply more than 3 pints (1.3 lbs. a.i./acre) of **Sequence** per acre per application.
- **DO NOT** apply more than 3 applications per year.
- **DO NOT** apply less than 10 days from the last application.
- **DO NOT** apply later than 5 weeks after petal fall.

APPLES

DISEASE	APPLICATION DIRECTIONS
Scab (<i>Venturia inaequalis</i>)	Use 1.5 pints per acre of Sequence beginning at green tip, and 7 days later to clean the orchard of overwintering scab at the start of the season. Alternatively, apply Sequence preventively at weekly intervals from green tip to petal fall. DO NOT apply more than 2 applications per year. For resistance management, it is advised to tank-mix Sequence with a fungicide including captan or mancozeb based formulation, at labeled rates. When tank-mixing Sequence with a fungicide including captan or mancozeb, fill the spray tank with clean water half of final desired volume. Start agitation. Slowly add Sequence at the specified rate. Once mixed, slowly add a fungicide including, captan or mancozeb based formulation, at the specified rate. Add water to desired final volume. Mix thoroughly and apply at once. Maintain sufficient agitation during mixing and spraying operations to ensure a uniform spray mixture.

RESTRICTIONS:

- **DO NOT** apply more than 1.5 pints (0.65 lb. a.i./acre) of **Sequence** per acre in any single application.
- **DO NOT** apply more than 3 pints (1.3 lbs. a.i./acre) of **Sequence** per acre per year.
- **DO NOT** apply more than 2 applications per year.
- **DO NOT** apply less than 7 days from the last application.
- **DO NOT** apply after petal fall.

APRICOTS[*], NECTARINES, AND PEACHES
[*Not registered for use by California]

DISEASE	APPLICATION DIRECTIONS	USE RATE
Leaf Curl (<i>Taphrina deformans</i>)	Use as a dormant or delayed dormant application. Additional applications may be made just before buds swell in the spring.	3 pints/A (1.3 lbs. a.i./acre)
Brown rot blossom blight (<i>Monilinia laxa</i> / <i>Monilinia fructicola</i>) (States West of the Mississippi River)	For disease suppression , use at pink bud. Repeat application at full bloom and again at petal fall.	
Scab (<i>Venturia carpophila anamorph</i> ; <i>Cladosporium carpophilum</i>)	Use at petal fall.	
Shot hole (<i>Wilsonomyces carpophilus</i>)	Use at dormant state, and repeat application at petal fall.	
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 9 pints of Sequence per acre per year (3.9 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence per acre per application. • DO NOT apply more than 3 applications per year. • DO NOT apply less than 7 days from the last application. • DO NOT apply after petal fall. 		

BANANAS[*]
[Not registered for use by California]

DISEASE	APPLICATION DIRECTIONS
Sigatoka	Air application: Use 0.85 to 1.5 pints per acre of Sequence as needed to maintain Sigatoka control in a mixture of 19 to 25 pints water/acre and 4 pints mineral oil/acre. Repeat application thereafter at 7- to 15-day interval. It is advised for subsequent cover sprays to use Sequence in rotation with other labeled fungicides that have a different mode of action.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 7.5 pints of product per acre per year (3.2 lbs. a.i./acre/year). • DO NOT apply more than 1.5 pints (0.65 lb. a.i./acre) of Sequence per acre per application. • DO NOT apply more than 5 applications per year. • DO NOT apply less than 7 days from the last application. 	

BRAZIL NUTS[*], BUTTERNUTS[*], CASHEWS[*], FILBERTS[*], HICKORY NUTS[*], MACADEMIA NUTS[*], AND WALNUTS
[*Not registered for use by California]

DISEASE	APPLICATION DIRECTIONS	USE RATE
Brown rot blossom blight (<i>Monilinia laxa</i> / <i>Monilinia fructicola</i>)	For disease suppression , use at pink bud, full bloom, and petal fall.	3 pints/A (1.3 lbs. a.i./acre)
Scab (<i>Venturia carpophila anamorph</i> , <i>Cladosporium carpophilum</i>)	Use at petal fall. Repeat application 2 and 5 weeks later.	
Shot hole (<i>Wilsonomyces carpophilus</i>)	Use at leaf emergency (bloom through petal fall). Repeat application 2 and 5 weeks later.	
Anthraxnose (<i>Gnomonia leptostyla</i>)	Use when first leaves unfold. Repeat application at 10- to 14-day intervals.	
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 9 pints of Sequence per acre per year (3.9 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence per acre per application. • DO NOT apply more than 3 applications per year. • DO NOT apply less than 10 days from the last application. • DO NOT apply within 7 days of harvest. 		

CHERRIES (SOUR AND SWEET)

DISEASE	APPLICATION DIRECTIONS
Leaf Spot (<i>Blumeriella jaapii anamorph, Coccomyces hiemalis</i>)	Use 1.5 - 3 pints of Sequence per acre at petal fall or when first leaves unfold. Repeat application thereafter at 7- to 10-day intervals. If conditions favor leaf spot, continue applications through harvest. For resistance management, it is advised to tank-mix with a fungicide including a captan based formulation, at labeled rate. Fill the spray tank with clean water at half of the final desired volume. Start agitation. Slowly add Sequence at the specified rate. Slowly add a fungicide including, captan, at the labeled rate. Add water to desired final volume. Mix thoroughly and apply at once. Maintain sufficient agitation during mixing and spraying operations to ensure a uniform spray mixture. Post-Harvest Applications to Trees: Apply 1.5 - 3 pints of Sequence per acre immediately after harvest and as needed.
Brown rot blossom blight (<i>Monilinia laxa/Monilinia fructicola</i>) (States West of the Mississippi River)	For disease suppression, use 3 pints of Sequence per acre at the time flower buds begin to open (white bud-popcorn bloom stages). Repeat application at full bloom and again at petal fall. Applications of Sequence to cherries when temperatures and humidity are excessively high may cause spotting to foliage or fruit.
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 12 pints of Sequence per acre per year (5.2 lbs. a.i./acre/year). • DO NOT apply more than 6 applications per year (2 applications at 3 pints of Sequence per acre (1.3 lbs. a.i./acre), and 4 applications at 1.5 pints of Sequence per acre (0.65 lb. a.i./acre)). • DO NOT apply less than 7 days from the last application. • DO NOT apply within 7 days of harvest. 	

PEANUTS[*]

[*Not registered for use by California]

DISEASE	APPLICATION DIRECTIONS
Early Leaf Spot (<i>Cercospora arachidicola</i>) and Late Leaf Spot (<i>Cercosporidium personatum</i>)	<p>Ground and Air Applications:</p> <p>Sequence alone - Use a maximum of 0.9 - 1.5 pints Sequence per acre in sufficient water to achieve thorough plant coverage. Begin with applications, first approximately 35 days after planting, and repeat at 10- to 14-day intervals.</p> <p>Sequence in tank-mix - When tank-mixing with white mold products including tebuconazole, dose rate can be reduced down to 0.8 pints of Sequence per acre. Tank-mix incompatibilities were observed with some tebuconazole based formulations. Therefore, we always advise to test the compatibility of the products used in combination with Sequence at small scale before spraying or to ask your local distributor about potential incompatibility. The tank-mix fungicide partner must be used at the labeled rate. Applications are made at 10- to 14-day intervals, beginning approximately 55 days after planting. DO NOT apply within 14 days of harvest.</p>
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 4.5 pints of Sequence per acre per year (1.95 lbs. a.i./acre/year). • DO NOT apply more than 1.5 pints (0.65 lb. a.i./acre) of Sequence per acre per application. • DO NOT apply more than 5 applications per year when using lower rates. • DO NOT apply less than 10 days from the last application. • DO NOT apply within 14 days of harvest. • DO NOT feed hay or vines (peanuts) to livestock. 	

PEARS

DISEASE	APPLICATION DIRECTIONS
Scab (<i>Venturia pirina</i>) and leaf spot (<i>Fabraea</i> spp.)	Protection Schedule – Ground and Air application: Use 3 pints per acre of Sequence at 7-day intervals to maintain scab control beginning at green tip. Fill the spray tank with clean water half of final desired volume. Start agitation. Slowly add Sequence at the specified rate. Add water to desired final volume. Mix thoroughly and apply at once. Maintain sufficient agitation during mixing and spraying operations to ensure a uniform spray mixture.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 9 pints of Sequence per acre per year (3.9 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence per acre per application. • DO NOT apply more than 3 applications per year. • DO NOT apply less than 7 days from the last application. • DO NOT apply within 7 days of harvest. 	

PECANS[*]

[*Not registered for use by California]

DISEASE	APPLICATION DIRECTIONS
Scab, liver spot, brown leaf spot, downy leaf spot, leaf blotch, downy mildew	Ground and Air application: Sequence alone - Use 3 pints of Sequence per acre in sufficient water to obtain good coverage of spray at pre- and post-pollination periods. Apply every 10 to 21 days or as needed as determined by rainfall, from bud break through shell hardening. Sequence in tank-mix - In case of tank-mix with triphenyltin hydroxide or fenbuconazole use 1.5 pints of Sequence per acre. Sequence has been used effectively on most pecan varieties. Severe foliage injury has been experienced on Barton, Moore, Shawnee and Van Deman in the Southeast. Foliar bronzing has sometimes been observed on other varieties, especially when tank-mixed with certain miticides.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 18 pints of Sequence per acre per year (7.8 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence per acre per application. • DO NOT apply more than 6 applications per year. • DO NOT apply less than 10 days from the last application. • DO NOT apply after shucks have started to open. 	

PLUMS[*], PLUMS-CHICKASAW[*], PLUMS-DAMSON[*], PLUMS-JAPANESE[*], PLUMCOTS[*], PRUNE PLUMS[*]

[*Not registered for use by California]

DISEASE	APPLICATION DIRECTIONS	USE RATE
Leaf Curl (<i>Taphrina deformans</i>)	Use as a dormant or delayed dormant application. Additional applications may be made just before buds swell in the spring.	3 pints/A (1.3 lbs. a.i./acre)
Brown rot blossom blight (<i>Monilinia laxa</i> / <i>Monilinia fructicola</i>) (States West of the Mississippi River)	For disease suppression , use at pink bud. Repeat application at full bloom and again at petal fall.	
Scab (<i>Venturia carpophila</i> anamorph, <i>Cladosporium carpophilum</i>)	Use at petal fall. Repeat application at 2 and 5 weeks after petal fall.	
Shot hole (<i>Wilsonomyces carpophilus</i>)	Use at dormant stage, and repeat application at petal fall.	
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 9 pints of Sequence per acre per year (3.9 lbs. a.i./acre/year). • DO NOT apply more than 3 pints (1.3 lbs. a.i./acre) of Sequence per acre per application. • DO NOT apply more than 3 applications per year. • DO NOT apply less than 7 days from the last application. • DO NOT apply within 7 days of harvest. 		

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container. **DO NOT** store in areas where temperatures are below 32 degrees F. Store in a cool, dry (preferably locked) area away from other pesticides, fertilizer, food, or feed that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Rigid Plastic Less Than or Equal to 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by state and local authorities.

Rigid Plastic Greater Than 5 Gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of empty container in a sanitary landfill or by other procedures allowed by state and local authorities.

Rigid Plastic Bulk and Mini-Bulk Containers: Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

DODINE GROUP U12 FUNGICIDE

Sequence

For control of certain diseases of Almonds, Apples, Apricots[*], Bananas[*], Beech nuts[*], Brazil nuts[*], Butternuts[*], Cashews[*], Cherries, Chestnuts[*], Chinquapins[*], Filberts[*], Hickory nuts[*], Macadamia nuts[*], Nectarines, Peaches, Peanuts[*], Pears, Pecans[*], Plums[*], Plums-chickasaw[*], Plums-damson[*], Plums-Japanese[*], Plumcots[*], Prune plums[*], and Walnuts.

[*Not registered for use by California.]

ACTIVE INGREDIENT:	WT. BY %
Dodine (n-dodecylguanidine acetate)	39.6%**
OTHER INGREDIENTS:	60.4%
TOTAL:	100.0%

**Equivalent to 3.4 pounds Dodine per gallon

KEEP OUT OF REACH OF CHILDREN DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

FIRST AID - IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to by a poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBERS - For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>. Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222.

NOTE TO PHYSICIAN - Probable mucosal damage may contraindicate the usage of gastric lavage.

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - DANGER/PELIGRO - Corrosive. Causes irreversible eye damage. Causes skin irritation. **DO NOT** get in eyes, on skin, or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid breathing spray mist or vapor. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. **ENVIRONMENTAL HAZARDS** - This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT**

contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **Surface Water and Erosion Control Statement:** This product may contaminate water through runoff and has the potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination. **DIRECTIONS FOR USE** - It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE:** Keep this product in its tightly closed original container. **DO NOT** store in areas where temperatures are below 32 degrees F. Store in a cool, dry (preferably locked) area away from other pesticides, fertilizer, food, or feed that is inaccessible to children and animals. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER HANDLING: Rigid Plastic Less Than or Equal to 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by state and local authorities. **Rigid Plastic Greater Than 5 Gallons:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of empty container in a sanitary landfill or by other procedures allowed by state and local authorities. **Rigid Plastic Bulk and Mini-Bulk Containers:** Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities. **CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!**

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

Manufactured For: Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707

EPA Reg. No. 82633-104-83529

EPA Est. No. (AG) 72159-GA-001; (MC) 89332-GA-001; (SC) 39578-TX-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: **2.5 Gals.*** **265 Gals.**

* Unless alternate checked

OPEN HERE