

DEAE Persefose BB

1. Basic product information

DEAE Persefose BB is a new type of highly cross-linked agarose chromatography resin formed by bonding diethylaminoethyl groups to high-flow agarose gel filtration chromatography resin. A weak anion exchange resin. It has high flow rate, low back pressure, high dynamic capacity, good chemical stability and mechanical properties, low non-specific adsorption, high recovery rate and convenient scale-up, shortening production time and improving production efficiency. It is widely used in the ion exchange separation of proteins, nucleic acids and polypeptides downstream of biopharmaceuticals and bioengineering.

2. Chromatography resin parameters

Resin type	Weak anion exchange
Matrix	Highly cross-linked agarose
Functional group	Diethylaminoethyl
Resin particle size	200 µm
Total ionic capacity	0.11–0.16 mmol Cl ⁻ /ml
Dynamic binding capacity	>50 mg BSA/ml
Recommended flow rate	1200–1800 cm/h
Maximum flow rate	1800 cm/h
Maximum working pressure	3 bar
Working temperature	4–40°C

3. Chemical resistance

pH stability*	2–12 (long term), 1–14 (short term, CIP)
Chemical stability	All commonly used aqueous buffers, 1M sodium hydroxide; 8M urea; 8M guanidine hydrochloride; 70% ethanol
Avoid	Oxidising agents, anionic detergents

* The physical and chemical properties and functions of the chromatographic resin have no obvious changes after being placed in an environment of 40°C and pH 2–12 after 7 days.

** v/v, volume ratio

4. Method of use

1. **Buffer selection:** the buffer salt whose buffer group does not interact with the chromatographic resin should be selected. If the binding and elution mode is used, the equilibration buffer should be a buffer with low salt (less than 5 mS/cm) and high pH (usually 1 pH unit higher than the isoelectric point of the target molecule) to facilitate the binding of the target molecule. Consider the stability of the sample in the buffer; the elution buffer is usually a buffer with a high concentration of salt (such as 1M NaCl) added to the equilibration buffer. If the flow-through mode is used, the equilibration buffer should adopt conditions that are conducive to the binding of impurities. After the target molecule has completely flowed through, it should be washed directly with buffer with high-concentration salt.
2. **Flow rate:** generally choose a linear flow rate of 100~600 cm/h according to the column bed height.
3. **Sample pretreatment:** to prevent the sample from clogging the column, the sample needs to be filtered with a 0.45 µm microporous membrane before loading it. It is recommended that the pH and conductivity of the sample is adjusted to be consistent with the equilibration buffer (dilution, ultrafiltration and desalting can be used to adjust the pH and conductivity of the sample).

4.2 Chromatography steps

1. **Equilibration:** use equilibration buffer to fully equilibrate the chromatography column until the pH and conductivity are stable and basically consistent with the equilibration buffer. This step usually requires 3–5 column bed volumes (CV).
2. **Sample loading*:** according to the binding capacity measured in the small test, determine the sample loading volume and loading amount of the sample on the DEAE Persefose BB.
3. **Washing*:** use equilibration buffer or other suitable buffer to wash the chromatography column until the UV stabilises and returns to the baseline.
4. **Elution*:** elution is achieved by increasing the concentration of salt ions. The concentration of salt ions in the elution buffer can be gradually increased through a linear gradient or a step gradient to elute molecules with different binding strengths. pH gradient elution or mixed elution can also be used.
5. **Regeneration:** rinse the column with a high-salt buffer (such as 2M NaCl).
6. **Re-equilibration:** re-equilibrate the column with equilibration buffer.

** If the flow-through mode is used, the 'sample loading' step should be set to collect; the 'washing' step should stop the collection after all the target molecules have passed through; the 'elution' step should directly use high salt buffer – impurities can be washed away.*

5. Cleaning and regeneration

Contaminants (e.g. lipids, endotoxins and proteins) accumulate on the column as the number of uses of the chromatography resin increases. Determine the frequency of CIP according to the degree of contamination of the chromatography resin (if the contamination is considerable, CIP is recommended after each use to ensure repeatability of results and to prolong the working life of the chromatography resin). For different types of impurities and contaminants, the recommended cleaning conditions are as follows:

- **Removal of strongly binding proteins:** wash with 5 CV of 2M NaCl solution, or use a high salt buffer not lower than pH 2, such as 1M NaAc solution.

- Removal of strongly hydrophobic proteins and precipitated proteins: first wash with 5 CV of 1M NaOH solution, then wash the lye with 5–10 CV of ultra pure or pure water.
- **Removal of lipoproteins and lipids:** first wash with 5 CV of 70% ethanol or 30% isopropanol, then rinse with 5–10 CV of ultra pure or pure water.

Note: 70% ethanol or 30% isopropanol should be degassed before use; the flow rate should be 30–60 cm/h during CIP. Reverse cleaning should be used when the clogging is severe.

To reduce the microbial load, it is recommended that 0.5~1M NaOH solution is used to treat the chromatographic resin. Treatment time is 15~30 minutes.

6. Storage

Keep the unopened chromatography resin in the original container and store at 4~30°C in a well-ventilated, dry and clean place. Do not freeze. Wash the used column with 2–3 CV of 20% ethanol solution and store at 2~8°C.

7. Destruction and recycling

Since chromatography resin is difficult to degrade in nature, it is recommended that the waste chromatography resin is incinerated, to protect the environment. For chromatography resin that has been in contact with biologically active samples such as viruses and blood, follow the local biosafety requirements before destroying or disposing of it.

8. Packing method

Detailed information on resin packaging is available on request. Please contact your local distributor.

9. Ordering information

Product name: DEAE Persefose BB

Product Code in Catalogue	Product Volume
296-00025	25 ml
296-00100	100 ml
296-00500	500 ml
296-01000	1 L
296-05000	5 L
296-10000	10 L

Document version: v1.0. Copyright © 2026 LT Biotech, UAB. All rights reserved. Any redistribution or reproduction of part or all the content in any form is prohibited without the express written permission of LT Biotech, UAB.