



# SERMORELIN 10 MG - VIAL

## RESEARCH USE PROTOCOL

Reconstitution	Reconstitute by adding 6 mL of bacteriostatic water to the vial.
Dosage	5 times per week (Monday–Friday) Draw 15 units (250 mcg)
Time of Day	PM (before bed), preferably on an empty stomach
Injection Type	Subcutaneous (abdomen, thigh, or upper arm)
Product Details	Concentration: 10 mg / 6 mL
Product Duration	One vial will last 2 months of dosing
Program Duration	2 months; cycle 1 week off between each month
Storage	Store refrigerated at 2–8°C (36–46°F). Do not freeze. Protect from light.

# WHAT IS SERMORELIN ?

Sermorelin is a peptide studied for its interaction with growth hormone-releasing hormone (GHRH) signaling pathways and endocrine system regulation.

It is commonly explored in research related to hormonal signaling, metabolic processes, and recovery-related pathways.

## WHAT'S IN THE BOX?



## HOW IT WORKS

### MECHANISM OF ACTION

Sermorelin is studied for its interaction with hormonal signaling pathways:

Associated with GHRH receptor-related signaling

Linked to growth hormone release pathways

Studied in IGF-1-related processes

Associated with metabolic regulation signaling

Connected to recovery and sleep-related pathways

These mechanisms are associated with endocrine signaling and metabolic processes.

## RESEARCH OBSERVATIONS

Studied for hormonal signaling pathways

Studied for metabolic regulation processes

Studied for recovery-related mechanisms

Studied for sleep-related signaling

Studied for body composition-related pathways





## OBSERVED REACTIONS IN RESEARCH SETTINGS

Research observations have noted mild and temporary responses such as injection site irritation, redness, mild headache, or increased hunger. Responses may vary depending on protocol design and individual variability.

## RESEARCH NOTES

In research settings, consistent dosing and timing of administration may influence observed outcomes, particularly in relation to sleep and hormonal signaling patterns. Individual variability should be considered when interpreting results.

## IMPORTANT CONSIDERATIONS FOR RESEARCH USE

---

Not intended for human consumption or therapeutic use

---

Not suitable for use during pregnancy or breastfeeding

---

Not recommended for individuals with severe medical conditions

---

Not recommended for individuals undergoing active medical treatment without supervision

---

Use in research settings may require professional oversight

---

Individual variability may influence observed outcomes