



BPC-157 / TB-500- 10MG / 10MG REBUILD - VIAL

RESEARCH USE PROTOCOL

Reconstitution	Reconstitute by adding 4 mL of bacteriostatic water to the vial.
Dosage	5 times per week (Monday–Friday) Draw 20 units (500 mcg)
Time of Day	PM (before bed), preferably on an empty stomach
Injection Type	Subcutaneous (abdomen, thigh, or upper arm)
Product Details	Concentration: 10 mg / 10 mg / 4 mL
Product Duration	One vial will last 1 month of dosing
Program Duration	2 months; cycle 1 week off between each month
Storage	Store refrigerated at 2–8°C (36–46°F). Do not freeze. Protect from light.

WHAT IS REBUILD ?

Rebuild is a peptide combination of BPC-157 and TB-500 studied for its role in tissue repair, inflammatory signaling, and recovery-related pathways.

It is commonly explored in research related to connective tissue support, recovery processes, and cellular repair mechanisms.

WHAT'S IN THE BOX?



HOW IT WORKS

MECHANISM OF ACTION

BPC-157 and TB-500 are studied for their complementary interaction with cellular repair pathways:

Associated with cellular repair and regeneration processes

Linked to angiogenesis (formation of new blood vessels)

Supports collagen-related signaling pathways

Associated with inflammatory response modulation

Supports cellular migration and recovery pathways

These mechanisms are associated with tissue repair and recovery processes.

RESEARCH OBSERVATIONS

Studied for tissue repair pathways

Studied for inflammation-related processes

Studied for recovery from physical stress

Studied for cellular regeneration mechanisms





OBSERVED REACTIONS IN RESEARCH SETTINGS

Research observations have noted mild and temporary responses such as localized irritation, redness, swelling, or mild fatigue. Responses may vary depending on protocol design and individual variability.

RESEARCH NOTES

Research observations have noted mild and temporary responses such as localized irritation, redness, swelling, or mild fatigue. Responses may vary depending on protocol design and individual variability.

IMPORTANT CONSIDERATIONS FOR RESEARCH USE

Not intended for human consumption or therapeutic use

Not suitable for use during pregnancy or breastfeeding

Not recommended for individuals with certain medical conditions

Use in research settings may require professional oversight

Not for use alongside medical treatments without supervision

Individual variability may influence observed outcomes