Bariatric Surgery Hydration Guide

YOUR HYDRATION GUIDE AFTER BARIATRIC SURGERY

The One Habit That Keeps You Out of the Hospital

Why this guide exists

Of everything we ask you to do after surgery, drinking enough fluid is the single most important, and the single most commonly missed.

Here is the reason. In a national study of 130,007 bariatric patients, the number one cause of being readmitted to the hospital in the first 30 days was not a leak. It was not bleeding. It was nausea, vomiting, and dehydration. That group alone accounted for more than a third of all readmissions.

In another national analysis, patients who needed treatment for dehydration were far more likely to end up in the emergency room than patients who stayed hydrated.

Almost none of that is caused by the surgery going wrong. Most of it is caused by simply not drinking enough.

The good news is that this is the most preventable complication you face.

Your daily target

At least 64 ounces of fluid every day. That is eight 8-ounce cups, or about 1.8 liters.

This is the number your surgical team will ask you about at every visit. Within several days of surgery you should be reliably hitting it.

Some patients need more, particularly in hot weather, with exercise, or if you are

© 2026 Nestor de la Cruz-Muñoz, MD. All rights reserved. No reproduction or distribution without written permission. https:www.delacruzmunoz.com

having any vomiting or loose stools.

Why this is harder than it sounds. Before surgery, you could drink 16 ounces in a few seconds. Now your stomach holds a few ounces at a time. You cannot drink your way to 64 ounces in three sittings anymore. You have to spread it across the entire day.

One study following patients through their first year after surgery found that oral fluid intake dropped by roughly half after surgery. That happens without patients even noticing. Which is exactly why this has to become a deliberate, scheduled habit rather than something you do when you feel thirsty.

Do not wait until you are thirsty

This is the most important mindset change in this guide.

After surgery, thirst is an unreliable signal. Many patients simply do not feel thirsty, and by the time you do, you are already behind.

Drink on a schedule, not on a feeling.

The five rules of drinking after surgery

1. Sip. Never gulp. Small, steady sips all day long. A gulp is too much volume too fast for your new stomach and will cause pain, nausea, or vomiting.

2. No straws. Straws pull air into your stomach, which causes gas, bloating, and pressure.

3. No carbonation. No soda, no sparkling water, no seltzer, no beer. Carbonation causes discomfort and pressure in your new stomach.

4. Never drink with meals. Stop drinking 30 minutes before you eat. Wait 30

© 2026 Nestor de la Cruz-Muñoz, MD. All rights reserved. No reproduction or distribution without written permission. https:www.delacruzmunoz.com

minutes after you finish before you drink again. Fluid taken with food washes the food through, so you get hungry sooner, and it fills space you need for protein.

5. Keep a bottle within arm's reach at all times. Bedside, car, desk, couch. If you have to get up to get a drink, you will not drink enough.

What counts toward your 64 ounces

Yes, these count:

- Water, plain or flavored with sugar-free additives

- Protein water and protein shakes

- Sugar-free, non-carbonated flavored waters

- Decaffeinated tea and decaffeinated coffee

- Sugar-free popsicles (one popsicle is roughly 2 ounces)

- Sugar-free gelatin

- Clear broth, low sodium

- Milk and unsweetened milk alternatives, once you are on full liquids

- Sugar-free electrolyte drinks

No, these do not:

- Anything carbonated

- Regular soda, juice, sweet tea, sports drinks with sugar, and sweetened coffee drinks. Sugar can trigger dumping syndrome, especially after gastric bypass.

© 2026 Nestor de la Cruz-Muñoz, MD. All rights reserved. No reproduction or distribution without written permission. https:www.delacruzmunoz.com

- Alcohol, which should be avoided for at least the first 6 months

- Anything you have to drink through a straw

A note on caffeine. Small amounts of caffeine are usually acceptable once you are further out, but caffeine is mildly dehydrating and can irritate a healing stomach. Ask us before adding it back, and never count a caffeinated drink as a full contribution to your daily total early on.

How to actually hit 64 ounces: a sample day

The goal is roughly 4 to 8 ounces per hour while you are awake.

Time	What to drink	Ounces
7 AM	Water on waking	4
8 AM	Protein shake, sipped over an hour	8
9 AM	Water	4
10 AM	Water or decaf tea	4
11 AM	Water	4
12 PM	MEAL, no fluids	0
1 PM	Water, starting 30 min after eating	4
2 PM	Protein shake or water	8
3 PM	Water	4
4 PM	Water	4
5 PM	MEAL, no fluids	0
6 PM	Water, starting 30 min after eating	4
7 PM	Water or broth	6
8 PM	Water	4
9 PM	Water, small amount	4
	TOTAL	**62 to 66**

Adjust the timing to your own schedule. The structure is what matters: something in

© 2026 Nestor de la Cruz-Muñoz, MD. All rights reserved. No reproduction or distribution without written permission. https:www.delacruzmunoz.com

your hand almost every hour, and a protected 30-minute window on either side of every meal.

Tricks that work

Use a marked bottle. A 32-ounce bottle with time markers means you only have to finish it twice. Refilling it becomes the goal instead of counting cups.

Set a timer. Every 15 to 20 minutes for the first few weeks. Most patients need the reminder for about a month before it becomes automatic.

Try different temperatures. Many patients tolerate room temperature or warm fluids far better than ice cold in the early weeks. Some find the opposite. Experiment.

Flavor it. Sugar-free flavor drops, a squeeze of lemon, cucumber, or mint. Plain water becomes unappealing to a lot of patients after surgery, and that is a common reason intake drops.

Sip warm broth in the evening. It counts as fluid, adds a little sodium, and feels like a meal when you are missing eating.

Freeze it. Sugar-free popsicles or ice chips work well on days when liquid feels heavy.

Track it for the first month. Use the log below, a phone app, or move a rubber band up your bottle with each refill. Bring the numbers to your first appointment.

When drinking is hard

Certain days will be difficult. Here is what to do.

If fluid feels like it is sitting in your chest: slow down and take smaller sips. Try

© 2026 Nestor de la Cruz-Muñoz, MD. All rights reserved. No reproduction or distribution without written permission. https:www.delacruzmunoz.com

sipping while sitting upright and walking a bit afterward.

If you feel nauseated: stop for 15 minutes, then restart with tiny sips of something cold or something warm, whichever you tolerate better. Do not push through and do not skip the rest of the day.

If water tastes strange or metallic: this is common. Switch to flavored sugar-free water, broth, or decaf tea until it passes.

If you are simply not thirsty: expected. Drink on the clock anyway.

If you are vomiting: call us. Do not wait to see if it passes.

Warning signs of dehydration

Learn these. They come on gradually and patients often miss them until they are quite far behind.

Early signs:

- Dark yellow urine, or urinating much less than usual

- Dry mouth, dry lips, sticky saliva

- Headache

- Fatigue beyond what you expected

- Dizziness when standing up

- Muscle cramps

- Feeling irritable or unable to concentrate

Call us right away if you have:

© 2026 Nestor de la Cruz-Muñoz, MD. All rights reserved. No reproduction or distribution without written permission. https:www.delacruzmunoz.com

- No urination for more than 8 hours, or urine that is very dark

- Dizziness or feeling like you may faint

- A resting heart rate over 100 beats per minute

- Vomiting that will not stop

- Inability to keep fluids down for 8 to 12 hours

- Confusion or unusual sleepiness

24-hour number: [PHONE]

Getting a bag of IV fluid in the office is a small thing. Being admitted with kidney injury is not. Call early.

A simple check: look at the color of your urine. Pale yellow or nearly clear means you are doing well. Dark yellow or amber means drink more today.

Hydration is not just a first-month issue

Most patients focus hard on fluid for a few weeks and then drift.

Your capacity stays limited for life. So does your need for 64 ounces a day. Dehydration remains one of the most common reasons bariatric patients seek urgent care years after surgery, particularly during illness, hot weather, travel, or any period of vomiting or diarrhea.

Be extra deliberate when you are:

- Sick with any stomach bug

- Traveling, especially by air

© 2026 Nestor de la Cruz-Muñoz, MD. All rights reserved. No reproduction or distribution without written permission. https:www.delacruzmunoz.com

- Exercising or outdoors in the heat

- Working a long shift without easy access to a bottle

14-DAY HYDRATION LOG

Bring this to your first postoperative appointment.

Day	Total ounces	Urine color (pale / dark)	Any vomiting?
1			
2			
3			
4			
5			
6			
7			
8			
9			
10			
11			
12			
13			
14			

THE SHORT VERSION

- **64 ounces every single day**

- **Sip constantly. Never gulp.**

© 2026 Nestor de la Cruz-Muñoz, MD. All rights reserved. No reproduction or distribution without written permission. https:www.delacruzmunoz.com

- **No straws. No carbonation. No sugar.**

- **Nothing to drink 30 minutes before or after meals**

- **Drink on a schedule, not when you feel thirsty**

- **Pale urine means you are on track**

- **Bottle within arm's reach, always**

Dehydration is the most common reason bariatric patients go back to the hospital. It is also the easiest one to prevent.

Real science. Real surgery. Real results.

© 2026 Nestor de la Cruz-Muñoz, MD. All rights reserved. No reproduction or distribution without written permission. https:www.delacruzmunoz.com